

Close-to-threshold Meson Production in Hadronic Interactions*

P. Moskal,^{1,2} M. Wolke,¹ A. Khoukaz,³ W. Oelert¹

¹IKP, Forschungszentrum Jülich, D – 52425 Jülich, Germany

²IP, Jagellonian University, PL – 30-056 Cracow, Poland

³IKP, Westfälische Wilhelms-Universität Münster, D – 48149 Münster, Germany

Abstract

Studies of meson production at threshold in the hadron-hadron interaction began in the fifties when sufficient energies of accelerated protons were available. A strong interdependence between developments in accelerator physics, detector performance and theoretical understanding led to a unique vivid field of physics. Early experiments performed with bubble chambers revealed already typical ingredients of threshold studies, which were superseded by more complete meson production investigations at the nucleon beam facilities TRIUMF, LAMPF, PSI, LEAR and SATURNE. Currently, with the advent of the new cooler rings as IUCF, CELSIUS and COSY the field is entering a new domain of precision and the next step of further progress.

The analysis of this new data in the short range limit permits a more fundamental consideration and a quantitative comparison of the production processes for different mesons in the few-body final states. The interpretation of the data take advantage of the fact that production reactions close-to-threshold are characterized by only a few degrees of freedom between a well defined combination of initial and exit channels. Deviations from predictions of phase-space controlled one-meson-exchange models are indications of new and exciting physics. Precision data on differential cross sections, isospin and spin observables — partly but by no means adequately available — are presently turning up on the horizon. There is work for the next years and excitement of the physics expected. Here we try to give a brief and at the same time comprehensive overview of this field of hadronic threshold production studies.

PACS numbers:

13.60.Hb, 13.60.Le, 13.75.Cs, 13.75.-n, 13.85.Lg, 13.85.Ni, 13.85.Rm,
14.40.Cs, 25.40.-h, 25.40.Ve, 29.20.Dh

*This review article will be published in “Prog. Part. Nucl. Phys.” Vol. 49, issue 1 (2002).

Contents

1	Introduction	4
1.1	Aspects of threshold production	6
1.2	Hadronic and electromagnetic probes	7
1.3	Hadronic interactions	9
2	Basics of free and quasi-free production	11
2.1	Free NN scattering	11
2.2	Scattering inside the deuteron	18
3	Phenomenology of the initial and final state interaction	23
4	Single meson production in NN scattering	38
4.1	Dynamics of pseudoscalar meson production	38
4.2	Vector meson production	47
4.3	Scalar sector	52
4.4	Associated strangeness production	55
5	Meson pair production	65
5.1	Double pion production in NN scattering	65
5.2	Double pion production in pd scattering	67
5.3	Double K-meson production in pd scattering	69
5.4	Strangeness dissociation into $K\bar{K}$ -pairs	70
6	Meson production on light nuclei	72
6.1	Dynamics in three and four nucleon systems	72
6.2	Quasi-bound states	75
6.3	Test of invariances	76
7	Experimental facilities	79
7.1	Accelerators for hadronic physics at medium energies	79
7.2	Targets	80
7.2.1	Fiber Targets	81
7.2.2	Liquid Targets	81
7.2.3	Pellet Targets	81
7.2.4	Gas Jet Targets	82
7.2.5	Cluster Targets	83
7.3	Typical detector arrangements	83
7.3.1	Example of a 4π internal-detector	84
7.3.2	Example of a centre-of-mass 4π external-detector	85
7.3.3	Example of a dedicated internal threshold detector	86
8	Outlook	88
8.1	Polarized beams and/or targets	88
8.2	Final state interaction	90
8.3	Baryon resonances	90
8.4	Symmetries	90
8.5	Meson production in the three-nucleon system	91
8.6	Apparative aspects	91

8.7	Epilogue	91
-----	--------------------	----

1 Introduction

The complexity of the hadronic structure is one of the actual and topical fields of physics concerning the microscopic scale. Different, complementary as well as competing experimental approaches are used to attack the challenge of the hadronic spectroscopy.

A detailed understanding of the strong interaction dynamics on the symmetries of QCD is a fundamental aim of hadron physics. At high energies the QCD serves as a reliable theory whereas at low energies the non-perturbative QCD phenomena must be investigated. Here calculations are performed in an effective field theory of hadrons represented by the chiral effective Lagrangian, a theory of Goldstone bosons coupled to the octet of baryons and vector mesons. The QCD Lagrange function is written as:

$$L_{QCD} = \bar{q} (i\gamma^\mu \partial_\mu - M) q + L_{gluon} + L_{quark-gluon}. \quad (1)$$

In principle, q describes the column vector of the quarks u, d, s, c, b, t with each of them represented by a spinor, identical to the Dirac equation for spin- $\frac{1}{2}$ electrons. At energies in the few GeV range only the first three quarks are of any relevance, M represents the diagonal of the matrix for the quark current masses.

Data of high quality and precision on hadronic processes at low and intermediate momenta are necessary in order to verify the systematic low energy expansion of the Chiral Perturbation Theory (χPT), which has already enforced an important insight into the structure and dynamics of nucleons and mesons [1, 2]. It is known that the current masses of the lightest three quarks are significantly smaller than the typical hadronic energy scale, represented by the proton mass of $m_p \approx 1 \text{ GeV}/c^2$. This is the reason why in first order the quark masses of the light quarks are neglected in theoretical considerations and the spin- $\frac{1}{2}$ fields can be separated into two independent left- and right-handed parts, reflected by the chiral symmetry. The QCD Lagrangian separates into two identical images:

$$L_{QCD}[\bar{q}, q] = L_{QCD}[\bar{q}_{left}, q_{left}] + L_{QCD}[\bar{q}_{right}, q_{right}]. \quad (2)$$

Left-handed quarks do not communicate with right-handed quarks and *vice versa*. There is no Lorentz transformation which can change the handedness of a mass-less quark or particle. This symmetry of the Lagrange function is not the symmetry of the spectrum of the particles since otherwise each hadron should have a partner of equal mass but inverted parity. Consequently there would be a second proton with negative parity which does not exist. There are no parity-doublets in the spectrum of the strong interacting particles and therefore the symmetry is spontaneously broken. Due to this breaking the mass-less so-called Goldstone bosons, the pseudoscalar π , K and η mesons appear. The restriction of the interaction of these particles among each other and with other hadrons can be used to analyze consequences of the chiral symmetry and its breaking in the framework of an effective field theory.

In reality the lightest hadrons — the Goldstone bosons — are by no means mass-less, which can be understood by an explicit breaking of the chiral symmetry including a term for the quark masses. For each massive Fermion there exists a Lorentz transformation which transforms a left-handed to a right-handed field. The effect of this explicit breaking now can be treated perturbatively since the masses of the light quarks are much smaller than the typical hadronic mass.

Predicting the low-energy properties of nuclear and particle physics, χPT plays an important role in understanding hadron physics in the non-perturbative regime. Therefore it is very important to know the properties of the mesons, their structure and interaction in the hadronic environment.

The physics program at the medium energy hadron accelerators was and is focusing on studies of the production and the decay of light mesons and baryon resonances and the conservation or violation of symmetries.

Following this sequence we will present and discuss production of mesons and meson pairs at threshold with the questions depicted by the interaction view of figure 1 [3] where in the nucleon–nucleon (NN) scattering a meson X is created in a one–boson–exchange model. For the particular case the questions have to be answered: how is the distortion of the incident NN waves (ISI) included, which mesons contribute to the exchange process, is there an intermediate baryon resonance, how significant are rescattering contributions of the exchange mesons and what is a reasonable treatment of the NN and NX final state interactions (FSI)? Especially for the NN –interaction the FSI is crucial because of the nearby poles in the S–wave amplitudes corresponding to the deuteron bound state in the 3S_1 channel or the 1S_0 virtual state [3]. These poles and the phase–space factors tend to determine much of the energy dependence of the total cross section for meson production. Furthermore, in any region where these poles dominate, it is possible to link quantitatively meson production in cases where the two nucleons emerge separately or as a bound deuteron state.




Figure 1: Diagrammatic view of the one–boson–exchange meson production process. Produced meson $= X$, nucleon momentum $= \vec{p}$, meson momentum $= \vec{q}$, relative NN momentum $= 2\vec{k}$ [3].

Even at threshold the reaction mechanism of the basic process for the interrelation between real pions and virtual exchange meson currents as for instance in the pion production ($NN \rightarrow NN\pi$) is still not fully understood. Though first data suggested that s–wave pions were produced in a heavy–meson–exchange process, later measurements resulted in an interpretation of an interference among transition amplitudes as Ss , Ps , Pp , Sd and Ds , where the capital letter indicates the NN final state wave and the small letter the angular momentum between the two nucleons and the meson produced. In addition, it has been concluded that for higher partial waves π exchange rather than a heavier meson exchange is more significant.

Especially the pion–nucleon interaction went through several stages of increasing sensitivity for tests of nuclear theory. As indicated above more than a decade after the initial work [4,5] only in the sixties explicit calculations on cross sections of the $pp \rightarrow pp\pi^0$, $pn\pi^+$ transitions were possible [6,7] which, however, failed the total cross section [8–10] by a factor of five. After further theoretical developments, see e.g. reference [11], only recently detailed calculations were provided within the Jülich meson exchange model [12–14] including transitions in the threshold region beyond $l_\pi = 0$ and predicting analyzing powers and spin correlation coefficients making use of i) the basic diagrams, ii) final state interactions, iii) off–shell effects, iv) the exchange of heavier mesons and v) influences of intermediate excitation of the Δ resonance. On the other hand, the Jülich model does not account for quark degrees of freedom i.e. calculations within the framework of the χPT which, in fact, predict cross sections too small [15]

compared to experiments.

The heavier the meson produced in nucleon–nucleon scattering the larger the momentum transfers and thus in such processes the short–range parts of the production operators are tested.

Double meson production processes with both mesons being either identical (e.g. $\pi^0\pi^0$ or $K^0\overline{K}^0$) or different (e.g. $\pi\eta$) are in principle similar to the single ones, however, the possibly associated baryon resonances as intermediate states differ significantly. For instance, the two–pion production very likely is dominated by either the $P_{11} N^*(1440)$ resonance — via an effective σ exchange giving information of a scalar meson excitation of the $P_{11} N^*(1440)$ resonance — or the simultaneously excited $P_{33} \Delta(1232)$ resonances. Due to selection rules here the choice of definite quantum numbers in the initial and final state might help to sort out the reaction mechanism.

In any case, a large amount of the knowledge about the π – π interaction has been obtained by the $\pi N \rightarrow \pi\pi N$ reaction, whereas data employing proton beams just start to come.

As long as only S–wave processes are involved, which is the privilege of threshold production studies, scattering length and effective range approaches are used to describe the interaction sign and strength. In case of strong attractive interactions a distinction between the final state scattering and the formation of a baryonic resonance leading to bound or quasi–bound states can not be made uniquely. Such investigation will be presented.

Isospin violation or charge symmetry breaking processes are a topical and interesting field of threshold production physics. The essential contribution to isospin breaking is the possible $\pi^0 - \eta$ mixing. Therefore experiments around the η threshold as e.g. $dd \rightarrow \alpha \pi^0(\eta)$ or $pd \rightarrow {}^3He \pi^0({}^3H \pi^+)$ should identify the $\pi^0 - \eta$ mixing angle which in turn might give hints to the mass difference of the up and down quarks.

1.1 Aspects of threshold production

Threshold production experiments are characterized by excess energies which are small compared to the produced masses ¹. Consequently, in fixed target experiments the momenta of the final state particles transverse with respect to the direction of the incoming beam are small compared to the longitudinal components. Thus, ejectiles are confined to a narrow forward cone in the laboratory system around the centre–of–mass movement and — close–to–threshold — an experimental acceptance covering the full solid angle is feasible with comparatively small dedicated detector arrangements.

Small relative momenta in the final state effectively limit the number of partial waves contributing, simplifying the theoretical interpretation of experimental results. It should be noted, that already three–body final states require — in principle — a three–body Faddeev like approach, which has not been accomplished so far [16,17]. However, as first described by Watson [18], with two strongly interacting particles in the final state, the energy dependence of the total cross section close–to–threshold is essentially determined by the (three–body) phase–space and the energy dependence of the on–shell final state interaction (FSI). Due to small relative velocities FSI effects are inherent to the experimental observables. Thus, the interpretation of data in terms of reaction dynamics requires a correct treatment of both initial ² (ISI) and final state interactions [21] (for a review see [22]). On the other hand, FSI effects might provide access to low–energy scattering parameters, which are difficult to obtain otherwise in case of unstable particles.

¹With meson production up to masses of $\approx 1 \text{ GeV}/c^2$ discussed in this article, we will principally refer to experimental results and theoretical investigations covering excess energies up to 140 MeV (the π –mass) in excess of the respective thresholds.

²At high relative momenta of the colliding nucleons necessary for meson production the on–shell nucleon–nucleon interaction exhibits a rather weak energy dependence and might well be approximated by an overall reduction of the cross section in magnitude [19,20]. However, for produced masses in the $1 \text{ GeV}/c^2$ regime, even for proton–proton scattering no reliable model exists to allow a consistent evaluation of the initial state interaction [20].

Meson production at threshold implies high momentum transfers Δp , given by

$$\Delta p = \frac{1}{2} \sqrt{(m_N + m_B + m_X)^2 - 4 m_N^2} \quad (3)$$

for the reaction type $NN \rightarrow NBX$ with N , B and X denoting a nucleon, baryon or meson in the initial and final state and m_N , m_B and m_X as the respective masses. With momentum transfers in the range of 0.37 GeV/c to 1.10 GeV/c for π^0 and ϕ meson production, threshold production probes with corresponding distances between 0.53 fm and 0.18 fm the short range part of the nucleon–nucleon interaction. Consequently, the energy dependence of the primary production amplitude is expected to be weak, motivating Watson’s approach [18].

Theoretical studies have been carried out mainly within the framework of hadronic meson exchange models, i.e. with baryons and mesons as effective degrees of freedom (for a recent review see [20]). Chiral perturbation theory has been applied for the description of data on π production close-to-threshold [23]. However, in view of the characteristic distances mentioned above, QCD inspired models [24, 25] with constituent quarks and gluons or instantons as relevant degrees of freedom, might turn out to be appropriate. High quality exclusive data on close-to-threshold meson production in the energy range of non-perturbative strong interaction physics will be crucial for exploring the boundary between effective meson exchange models and (so far phenomenological) approaches based on quark–gluon degrees of freedom.

1.2 Hadronic and electromagnetic probes

It seems that the χPT is of limited value once strangeness is involved in the hadronic systems and non-perturbative coupled-channel considerations beyond χPT are on the market [26, 27]. High quality data on the meson ($\pi, \eta, \eta', \omega, K^+, K^-$) production at threshold have been and are being produced at the hadronic beam cooler rings using the baryon number $B = 2$ systems (essentially pp and pn). At least equally important are studies with hadronic interactions in the $B = 1$ sector. However, most measurements with π - or K -beams used bubble chamber techniques which naturally suffer from very poor statistics.

Accurate investigations of differential cross sections for meson production from the initial π plus proton or K plus proton systems are needed at several momenta to study the s- and p-wave dynamics and their interplay. There are several indications that as intermediate states both the S_{11} resonance and the $K\Sigma$ quasi-bound state — followed by a final state interaction leading to nucleon and meson — are essential and could be investigated best via the $B = 1$ sector.

Until now, most of the data for threshold meson production were observed with proton beams. The use of antiprotons has contributed considerably to the knowledge of the meson spectroscopy up to serious candidates for glueballs and exotic hybrids but is still suffering from the lack of unique interpretations.

A comparison of the meson production in the fundamental process to productions on heavier nucleons would give constraints to eliminate uncertainties in the basic interactions. As outlined by W. Weise and U.-G. Meißner [28], precise measurements of such processes set the necessary constraints for the effective Lagrangian of low-energy QCD with strange quarks. Its detailed knowledge has impacts on several other important issues, such as the strange quark content of the nucleon.




Figure 2: Proton–proton cross sections for different reactions of the light meson production in the threshold region. For comparison also the total and elastic cross sections of proton–proton scattering are shown. Relevant references to the data will be given throughout the present article at the appropriate sections.

Though in the present article we will restrict ourselves to hadron induced reactions, in general induced with p^- , d^- , π^- and $\bar{p}$ -beams, investigations on the structure of the nucleon as seen with leptons should at least be mentioned [29]. The use of leptonic interactions via photon or electroweak boson exchanges has the attraction of combining a weakly interacting particle of well controlled energy momentum transfer with the complex multi particle system containing charged constituents. Photo-production provides a crucial complementary information compared to hadronic processes, since they are characterized by the electromagnetic probe as a virtual vector meson or an $s\bar{s}$ fluctuation and as such are a powerful source of spin–1 states. The well proven perturbative methods allow us to define a hierarchy of operators which, at least in principle, can be extracted separately. Perturbative methods finally allow to overcome one of the limitations of the electroweak interaction, that is the access to gluon distributions. In fact the strong interaction in the limit of hard scales is becoming a probe for the soft

region. All this can be complemented by the addition of the spin and flavour degrees of freedom [29].

Cross sections are very small in the threshold region down to eight orders of magnitude compared to the total yield. Due to the rapid growth of the phase-space volume the total cross section of the meson production reactions increases by orders of magnitudes in a few MeV range of excess energy. Such studies have been made possible only due to the low emittance and small momentum spread proton beams available at the storage ring facilities. Figure 2 demonstrates threshold production measurements of light mesons in the proton–proton interaction.

Especially hadron accelerators offer the possibility to excite nucleonic resonances as N^* ’s and $\Delta^{(*)}$ ’s with the same quantum numbers as those of the proton via the $p\alpha \rightarrow N^*\alpha$ reaction, where the α –channel — with its spin and isospin zero — serves as a spin–isospin filter.

This scalar–isoscalar tool is complementary to the vector excitation of nuclei with electromagnetic probes. Recent results from different electron beam facilities as ELSA and MAMI in Germany and NIKHEF in The Netherlands certainly deserve a detailed and circumstantial presentation. Due to the present limitation to hadronic interactions we can not give appropriate credit to this field of physics and refer to [29,30]. At the Continuous Electron Beam Accelerator (CEBAF) of the Thomas Jefferson National Accelerator Facility (TJNAF) an electron beam is used for exciting particular nuclear resonances and observing their individual decay modes.

1.3 Hadronic interactions

The interaction of hadrons — being the reflection of the strong force acting between their constituents — delivers indirect information about their structure and the strong interaction itself. In the frame of the optical potential model the hadronic interaction can be expressed in terms of phase-shifts, which in turn are described by the scattering length and effective range parameters. These are quite well known for the low-energy nucleon–nucleon interaction [31,32] yet they are still poorly established in case of meson–nucleon or even meson–meson interactions. This is partly due to the absorption of mesons when scattering on a baryon. To account for this effect the scattering length becomes a complex quantity where the imaginary part — e.g. in case of the nucleon– η interaction — describes the $\eta N \rightarrow \pi N$ and $\eta N \rightarrow \text{multi-}\pi N$ processes. Moreover, the short life-time of all *neutral* ground state mesons prohibits their utilization as secondary beams and therefore the study of their interaction with hadrons is accessible only via observations of their influence on the cross section of the reactions in which they were produced ($NN \rightarrow NN \text{ Meson}$). When created close to the kinematical threshold with the relative kinetic energy being in the order of a few MeV, the final state particles remain much longer in the range of the strong interaction than the typical life-time of N^* or Δ baryon resonances with 10^{-23} s and hence they can easily experience a mutual interaction before escaping the area of an influence of the hadronic force. This — as introduced by Watson [18] — final state interaction (FSI) may significantly modify both the original distributions of relative momenta of the outgoing reaction products and the magnitude of the production cross section. Though it is easy to understand that the FSI changes the distributions of the differential cross sections it is rather difficult to cope for the influences on the magnitude of the total reaction rate, since one tends to separate the primary production from the final state interaction in space as well as in time [18]. Considering the primary production as a separate process it is well worth trying to understand the phenomenon qualitatively. If there were no final state interactions the total cross section would be fully determined by the kinematically available phase-space volume, V_{ps} , where each interval is populated with a probability governed by the primary production amplitude only:

$$\sigma = \frac{1}{F} \int dV_{ps} |M|^2 \approx \text{const.} \cdot V_{ps}. \quad (4)$$

The approximation in the equation results from the assumption that $|M|^2 = \text{constant}$ in a few MeV range above the production threshold [33–35]. F denotes the flux factor of the colliding particles.

In the classical picture we might imagine that the reaction particles are created together with their appropriate force field and when escaping the interaction region they acquire a potential energy increasing or decreasing their kinetic energy depending whether the interaction is repulsive or attractive. For an attractive interaction they could be created also in those phase-space partitions which are not available for non-interacting particles and subsequently be “pulled down” to the energetically allowed regions by final state interaction. The temporary growth of the primary production phase-space would then increase the reaction rate. Contrary, in case of a repulsive interaction the particles must be produced in the lower phase-space volume since leaving the interaction area they will acquire additional kinetic energy. The reduction of the total cross section, for example in case of the repulsive Coulomb force, can easily be understood when considering the production in a coordinate space of point-like objects. Here — in contrast to non-interactive particles — a strongly repulsive object can not be produced at appropriately small distances since their later acceleration would lead to the breaking of the energy conservation and thus the space available to primary production is reduced. One can also argue relativistically that the primary mechanism creates the particles off the mass shell and subsequently they are lifted onto the mass shell by the final state interaction. The solid line in figure 3 (left) depicts the boundary of the Dalitz plot in case of the $pp \rightarrow pp\eta$ reaction calculated at the total centre-of-mass energy $\sqrt{s} = 2433.8 \text{ MeV}$ exceeding by 10 MeV the threshold energy. The area surrounded by that curve is a direct measure of the kinematically available phase-space volume. The dotted line shows the




Figure 3: (left) The solid line indicates the kinematically available area for the $pp\eta$ system with a total centre-of-mass energy $\sqrt{s} = 2433.8 \text{ MeV}$. The dotted line depicts the range assuming a reduction of the proton and η -meson masses by 2 MeV. The phase-space volume results from an integral over the closed area: $V_{ps} = \frac{\pi^2}{4s} \int \int d m_{p_1\eta}^2 d m_{p_2\eta}^2$. (The powers of 2π are skipped here and will be included into the flux factor F according to the convention introduced by Byckling and Kajantie [36]). (right) Distribution of the phase-space for the $pp\eta$ system modified by the proton-proton interaction and calculated for an excess energy of $Q = 16 \text{ MeV}$. The area of squares are proportional to the number of entries. The largest square of the figure corresponds to 260 events.

corresponding plot at the moment of the primary creation if the mass of each particle was reduced by 2 MeV, demonstrating that now the available phase-space grows significantly. Indeed, as shall be inferred from the experimental results presented in subsequent sections, at excess energies of a few MeV

above threshold, the mutual interaction among the outgoing particles enhances drastically — by more than an order of magnitude — the total cross section and modifies appreciably the occupation of the phase-space. Figure 3 (right) indicates the phase-space distribution expected for the $pp\eta$ system at an excess energy of $Q = \sqrt{s} - 2m_p - m_\eta = 16$ MeV, assuming a homogeneous primary production and taking into account the S-wave interaction between the protons. The proton–proton FSI modifies the homogeneous Dalitz plot distribution of “non-interacting particles”, enhancing its population at a region where the protons have small relative momenta. The interaction of the proton– η system would manifest itself at low invariant masses $m_{p\eta}^2$ corresponding to small relative momenta between the proton and the η meson. Such effects observed in the experiments are presented in section 4.

At the vicinity of the threshold, relative angular momenta larger than $l = 0 \hbar$ will not play any role due to the short range of the strong interaction and small relative momenta of the produced particles. It can be inferred from parity and angular momentum conservation laws that the production of an N – N – $Meson$ system (for pseudoscalar or vector mesons) in relative S-waves may only occur if the nucleons collide in P-wave. Thus at threshold the transition $P \rightarrow Ss$ is the only possible one, with capital letters denoting the angular momentum between nucleons and the small letter for the meson angular momentum with respect to the pair of nucleons. An interaction of the nucleons in the entrance channel influences the production process appreciably [21], similarly as that described above between the outgoing particles. For example, for η production in the $pp \rightarrow pp\eta$ reaction the initial state interaction (ISI) reduces the total cross section by about a factor of 3–5 [19, 21] due to the repulsive proton–proton 3P_0 -wave potential. This factor keeps constant in the range of a few tens of MeV [19] and hence, does not influence the energy dependence of the total cross section of the meson production, which remains predominantly determined by the final state interaction. In particular, the dominant S-wave nucleon–nucleon final state interaction is by far stronger than any of the low-energy meson–nucleon ones, with the only exception of the proton– K^- FSI (see section 3). The effect of the nucleon–nucleon FSI diminishes with increasing excess energy since it significantly influences only that partition of the phase-space at which the nucleons have small relative momenta. Whereas this fraction keeps constant, the full volume of the phase-space grows rapidly: An increase of the excess energy from $Q = 0.5$ MeV to $Q = 30$ MeV corresponds to a growth of V_{ps} by more than three orders of magnitude [37]. As a result the S-wave nucleon–nucleon FSI is of less importance for higher excess energies where it affects a small fraction of the available phase-space volume only. A more quantitative discussion about the influence of the ISI and FSI effects at close-to-threshold production cross sections will be presented in section 3.

2 Basics of free and quasi-free production

2.1 Free NN scattering

Investigations of the production of mesons and their interactions with nucleons are based on measurements determining the total and differential production cross sections and their dependence on the energy of the interacting nucleons. Therefore, to enable a quantitative discussion on the mechanisms leading to the transformation of the energy-of-motion of nucleons into matter in the form of mesons let us recall the formula of the reaction cross section. In case of the $NN \rightarrow NN Meson$ process — with the four-momenta of the colliding nucleons denoted by $\mathbb{P}_a$ and $\mathbb{P}_b$ and with $n = 3$ particles in the exit channel — this reads:

$$\sigma = \frac{1}{F} \int dV_{ps} |M_{ab \rightarrow 123}|^2 = \frac{1}{F} \int \prod_{i=1}^n d^4\mathbb{P}_i \cdot \delta(\mathbb{P}_i^2 = m_i^2) \cdot \Theta(\mathbb{P}_i^2) \cdot \delta^4(\mathbb{P}_a + \mathbb{P}_b - \sum_{j=1}^n \mathbb{P}_j) \cdot |M_{ab \rightarrow 123}|^2, \quad (5)$$

where $|M_{ab \rightarrow 123}|^2$ denotes the square of the Lorentz-invariant spin averaged matrix element describing the probability to create two nucleons and a meson with four-momenta of $\mathbb{P}_i = (E_i, \vec{p}_i)$ and $i = 1..n$,

respectively. The energy and momentum conservation as well as the on-shellness of the created particles is ensured by the Dirac- δ and the Heaviside- Θ functions. The formula holds also for $n \neq 3$.

The total cross section is then defined as an integral of the probabilities to populate a given phase-space interval over the whole kinematically available range — determined by energy and momentum conservation — normalized to the flux factor F of the colliding nucleons. In the case of non-interacting final state particles the matrix element close to threshold $|M_{ab \rightarrow 123}|^2$, is nearly constant [33–35] and hence the allowed phase-space volume, V_{ps} , is the decisive quantity which governs the growth of the cross section with increasing excess energy Q . The latter — defined as the total kinetic energy — is shared among the outgoing particles in the reaction centre-of-mass frame ³:

$$Q = \sqrt{s} - \sum_{i=1}^n m_i, \quad (6)$$

where $s = |\mathbb{P}_a + \mathbb{P}_b|^2 = |\sum_{i=1}^n \mathbb{P}_i|^2$ denotes the square of total centre-of-mass energy. Exactly at threshold, where the particles' kinetic energy in centre-of-mass system is equal to zero ($Q = 0$ MeV), the total reaction energy $\sqrt{s}$ amounts to: $\sqrt{s_{th}} = \sum_{i=1}^n m_i$. Before writing explicitly the formula for V_{ps} let us introduce the kinematical triangle function λ defined as [36]:

$$\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2yz - 2zx, \quad (7)$$

which enables the expression of many useful kinematical variables in a very compact and Lorentz-invariant form. In particular, the momenta of particle i and j in their rest system equal:

$$p_i^* = p_j^* = \frac{\sqrt{\lambda(s_{ij}, m_i^2, m_j^2)}}{2\sqrt{s_{ij}}}, \quad (8)$$

where $s_{ij} = |\mathbb{P}_i + \mathbb{P}_j|^2$ stands for the square of the invariant mass of the ij system considered as one quasi-particle. The above relation gives an expression for the flux factor F from equation (5) in terms of the colliding masses of the nucleons and the total energy s only, namely:

$$F = 4\sqrt{s} (2\pi)^{3n-4} p_a^* = 2 (2\pi)^{3n-4} \sqrt{\lambda(s, m_a^2, m_b^2)}, \quad (9)$$

where we have chosen the convention introduced by Byckling and Kajantie [36] and included the (2π) factors for the phase-space $(2\pi)^{3n}$ and for the matrix element $(2\pi)^{-4}$ into the definition of F . It is important to note that at threshold, for an excess energy range of a few tens of MeV, the small fractional changes of total energy ($\frac{Q}{\sqrt{s}} = \frac{Q}{m_1 + m_2 + m_3 + Q}$) causes weak variations of the flux factor and influences only slightly the shape of the energy dependence of the total cross section.

For a two-particle final state $ab \rightarrow 12$ (for instance for the reaction $pn \rightarrow d\eta$) the phase-space integral defined in equation (5) reduces to $V_{ps} := \int dV_{ps} = \frac{\pi}{\sqrt{s}} p_1^* = \frac{\pi}{2s} \sqrt{\lambda(s, m_1^2, m_2^2)}$ and the total cross section for such reactions (when neglecting variations due to the dynamical effects ($|M_{ab \rightarrow 12}| = \text{const.}$)) should increase linearly with the centre-of-mass momentum of the produced meson in the vicinity of the threshold. The total cross section can be expressed analytically as a function of the masses of the particles participating in the reaction and the square of total reaction energy s :

$$\sigma_{ab \rightarrow 12} = \text{const} \cdot \frac{V_{ps}}{F} = \frac{\text{const} p_1^*}{16\pi s p_a^*} = \frac{1}{16\pi s} \frac{\sqrt{\lambda(s, m_1^2, m_2^2)}}{\sqrt{\lambda(s, m_a^2, m_b^2)}}. \quad (10)$$

³Traditionally by centre-of-mass system we understand a frame in which the momenta of all particles add to zero, called sometimes more explicitly centre-of-momentum frame.

Near threshold, at a given excess energy $Q = \sqrt{s} - m_1 - m_2$, the emission of the reaction products in the centre-of-mass frame is isotropic and the whole dynamics of the process manifests itself in the absolute value of the transition matrix element $|M_{ab \rightarrow 12}|$. The underlying production mechanisms can also be extracted from the deviations of the total cross section energy dependence following the prediction of relation (10). A visualization of possible differences in the dynamics for the production of various mesons can be realized by comparing the total cross section of the studied reactions at the same value of the phase-space volume normalized to the flux factor (V_{ps}/F). It is important to recognize that the comparison at the same centre-of-mass meson momentum (p_2^*) gives direct information about the amplitude differences only if the production of mesons with the same masses is concerned (for example $pp \rightarrow d\pi^+$ and $nn \rightarrow d\pi^-$), yet the reactions $pn \rightarrow d\eta$ and $pn \rightarrow d\pi^0$ have by far different V_{ps} at the corresponding p_2^* .

However, if one is interested in the decomposition of the production amplitude according to the angular momenta of the final state particles then indeed the appropriate variable for the comparison even for particles with different masses is the meson momentum in the reaction centre-of-mass frame. Correspondingly for the more than two-body final state, the adequate variable is the maximum meson momentum, since it is directly connected to the maximum angular momentum by the interaction range. Considering for a three-body exit channel ($ab \rightarrow 123$) that the meson (m_3) possesses maximum momentum when the remaining two particles are at rest relative to each other and employing definition (8) one obtains:

$$q_{max} = \frac{\sqrt{\lambda(s, (m_1 + m_2)^2, m_3^2)}}{2\sqrt{s}}. \quad (11)$$

Contrary to the two-body final state, at a fixed excess energy the dynamics of the meson production associated with two or more particles reflects itself not only in the absolute value of the square of the matrix element but also in distributions of variables determining the final state kinematics. Usually, in non-relativistic calculations of the total cross section, one takes the Jacobi momenta, choosing as independent variables the 'q'-meson momentum in the reaction centre-of-mass frame and 'k'-momentum of either nucleon in the rest frame of the nucleon-nucleon subsystem. By means of the λ function they can be expressed as:

$$q = \frac{\sqrt{\lambda(s, s_{12}, m_3^2)}}{2\sqrt{s}}, \quad \text{and} \quad k = \frac{\sqrt{\lambda(s_{12}, m_1^2, m_2^2)}}{2\sqrt{s_{12}}}, \quad (12)$$

with $s_{12} = |\mathbb{P}_1 + \mathbb{P}_2|^2$ denoting the square of the invariant mass of the nucleon-nucleon subsystem. In a non-relativistic approximation the expression of the total cross section defined by formula (5) for a meson production reaction in nucleon-nucleon interactions of the type $NN \rightarrow NN \text{ Meson}$ simplifies to:

$$\sigma \propto \int_0^{q_{max}} k q^2 |M_{ab \rightarrow 123}|^2 dq. \quad (13)$$

Denoting by L and l the relative angular momentum of the nucleon-nucleon pair and of the meson relative to the NN system, respectively and approximating the final state particles by non-distorted plane waves (case of non-interacting objects), whose radial parts $\psi_l(q, r)$ are given by the spherical Bessel functions:

$$\psi_l(q, r) \propto j_l(qr) \xrightarrow{qr \rightarrow 0} \frac{(qr)^l}{(2l+1)!}, \quad (14)$$

an expansion of the transition amplitude $|M_{Ll}|^2$ for the Ll partial wave around $q = 0$ leads to

$$|M_{Ll}|^2 \propto q^{2l} k^{2L}. \quad (15)$$

Applying the above proportionality in equation (13) and solving the integral, yields the partial cross section

$$\sigma_{Ll} \propto q_{max}^{2L+2l+4} \propto \eta_M^{2L+2l+4}, \quad (16)$$

where $\eta_M = q_{max}/m_M$ with m_M denoting the mass of the created meson⁴. Thus — at threshold — for the Ss partial wave the cross section for the $NN \rightarrow NN \text{ Meson}$ reaction should increase with energy of the fourth power of η_M . The dimensionless parameter η_M was introduced by Rosenfeld [5] as a variable allowing for the qualitative estimations of the final state partial waves involved in pion production. He argued that if the phenomenon of pion production takes place at a characteristic distance R from the centre of the collision, with R in the order of $\hbar/Mc$, then the angular momentum of the produced meson is equal to $l = Rq = \hbar q/Mc$. Hence, η_M denotes the classically calculated angular maximum momentum of the meson relative to the centre-of-reaction. The same arguments one finds in the work of Gell-Mann and Watson [4], where the authors do not expect the range of interaction to be greater than the Compton wave-length of the produced meson ($\hbar/Mc$). However, it is rather a momentum transfer Δp between the colliding nucleons which determines the distance to the centre of the collision $R \approx \hbar/\Delta p$ at which the production occurs. Based on the data indications, it is emphasized in the original article of Rosenfeld [5] that R is slightly less than $\hbar/(2Mc)$, which is numerically close to the value of $\hbar/\Delta p$. Directly at threshold, where all ejectiles are at rest in the centre-of-mass frame, Δp is equal to the centre-of-mass momentum of the interacting nucleons and hence, exploring equation (8) it can be expressed as:

$$\Delta p_{th} = \frac{\sqrt{\lambda(s_{th}, m_a^2, m_b^2)}}{2\sqrt{s_{th}}} \xrightarrow{\text{for } pp \rightarrow ppX} \sqrt{m_p m_X + \frac{m_X^2}{4}}. \quad (17)$$

Though the present considerations are constrained to the spin averaged production only, it is worth noting that very close-to-threshold — due to the conservation laws and the Pauli excluding principle⁵ — for many reactions there is only one possible angular momentum and spin orientation for the incoming and outgoing particles.

Table 1: Partial wave transitions for the $pp \rightarrow pp \text{ Meson}$ and $nn \rightarrow nn \text{ Meson}$ reactions at threshold

type	meson	spin and parity	transition
pseudoscalar	π, η, η'	0^-	$^3P_0 \rightarrow ^1S_0 s$
vector	ρ, ω, ϕ	1^-	$^3P_1 \rightarrow ^1S_0 s$
scalar	a_0, f_0	0^+	$^1S_0 \rightarrow ^1S_0 s$

For example the production of neutral mesons with negative parity — as pseudoscalar or vector mesons — may proceed in the proton-proton collision near threshold only via the transition between

⁴In former works [4, 5] dealing only with pions this parameter is denoted by η , here in order to avoid ambiguities with the abbreviation for the eta-meson, we introduce an additional suffix M .

⁵The Pauli principle for the nucleon-nucleon system implies that $(-1)^{L+S+T} = -1$, where L , S and T denote angular momentum, spin and isospin of the nucleon pair, respectively. For example if the nucleon-nucleon wave function is symmetric in space ($L = 0$) and spin ($S = 1$) then it must be antisymmetric in isospin-space ($T = 0$) to be totally antisymmetric.

^3P and $^1\text{S}_0$ s partial waves (see table 1). Therefore only the collision of protons with relative angular momentum equal to $1\hbar$ and a parallel spin orientation — which flips during the reaction — may lead to the production of negative parity neutral mesons. In the case of the production of neutral scalar mesons, the protons or neutrons must collide with anti-parallel spin orientations which remains unchanged after the reaction. These simple considerations imply that in the close to threshold measurements with the polarized beam and target one should see a drastic effect in the reaction yield depending whether the polarization of reacting protons is parallel or anti-parallel. In the ideal case, having the product of beam and target polarization equal to unity, the close to threshold π^0 , η or η' production can be realized only in case of the parallel spin orientation of the reacting protons and for the anti-parallel polarizations zero yield should be observed. Indeed, in reality, strong differences in the yield for various combinations of beam and target polarization have been determined in the pioneering measurements of the reaction $\vec{p}\vec{p} \rightarrow pp\pi^0$ at the IUCF facility [38].

Due to both the strong nucleon–nucleon low–energy interaction in the $^1\text{S}_0$ state and the meson–nucleon forces in the exit channel, the assumption of the non–interactive plane waves leading to equation (16) failed when confronted with experimental data. However, these deviations give the possibility for determining the still poorly known nucleon–meson interactions.




Figure 4: Monte-Carlo Simulations: (a) Phase-space distribution for the $pp \rightarrow pp\eta$ reaction at $Q = 16$ MeV modified by the proton-proton final state interaction. (b) The dotted line shows the projection of the pure phase-space density distribution onto the s_{pp} axis and the solid curve presents its modification by the proton-proton FSI. (c) The square of the scattering amplitude for the $pp \rightarrow pp$ elastic scattering as a function of the proton-proton invariant mass in the range $2m_p < \sqrt{s_{pp}} < 2m_p + Q$, calculated according to formula (34) of section 3. (d) Phase-space density distribution modified by the proton- η interaction, with scattering length equal to $a_{p\eta} = 0.7 \text{ fm} + i 0.3 \text{ fm}$. The proton- η scattering amplitude has been calculated according to the equation (37). A detailed discussion of the nucleon-nucleon and nucleon-meson interaction will be presented in section 3.1.

The interaction between particles depends on their relative momenta. Consequently, for investigations of final state interactions more instructive coordinates, than the q and k momenta, are the squared invariant masses of the two-body subsystems [39]. These are the coordinates of the Dalitz plot. In the original paper [40] Dalitz has proposed a representation for the energy partitions of three bodies in an equilateral triangle whose sides are the axes of the centre-of-mass energies. He took advantage of the fact that the sum of distances from a point within the triangle to its sides is a constant equal to the height. Therefore, the height of the triangle measures the total energy $\sqrt{s} = E_1^* + E_2^* + E_3^*$ and interior points — fulfilling four-momentum conservation — represent energy partitions. For a constant $\sqrt{s}$ due to the energy conservation, without losing any information, it is enough to consider the projection on any of the $E_i E_j$ planes. The linear relation between E_i^* and s_{jk} ($s_{jk} = s + m_i^2 - 2\sqrt{s}E_i^*$) allows to use $s_{jk} s_{ik}$ or $E_i E_j$ coordinates equivalently, with the following relation between the phase-space intervals: $dE_i^* dE_j^* = \frac{1}{4s} ds_{jk} ds_{ki}$. According to the Kilian's geometrical representation [39] a Dalitz lies on a plane in the three dimensional space (s_{12}, s_{13}, s_{23}) orthogonal to the space diagonal. The plane including a

Dalitz plot corresponding to a given total energy s is then given by the following scalar product [39]:

$$(1, 1, 1)(s_{12}, s_{13}, s_{23}) = s_{12} + s_{13} + s_{23} = s + m_1^2 + m_2^2 + m_3^2. \quad (18)$$

The second equality of equation (18) means that there are only two independent invariant masses of the three subsystems and therefore a projection onto any of the $s_{ij} s_{jk}$ planes still comprises the whole principally accessible information about the final state interaction of the three-particle system. In the case of no dynamics whatsoever and the absence of any final state interaction the occupation of the Dalitz plot would be fully homogeneous since each phase-space interval would be equally probable. The final state interaction would then appear as a structure on that area. Figure 4a shows — for the example of the $pp \rightarrow pp\eta$ reaction — how the uniformly populated phase-space density is modified by the S-wave (1S_0) interaction between outgoing protons. An enhancement in the range corresponding to the low relative momentum between protons is clearly visible. A steep decrease of the occupation density with increasing invariant mass of the proton-proton subsystem is even better seen in figure 4b presenting the projection of the phase-space distribution onto the s_{pp} axis indicated by an arrow in figure 4a. This is a direct reflection of the shape of the proton-proton (1S_0) partial wave amplitude shown in figure 4c. Figure 4d shows the Dalitz plot distribution simulated when switching off the proton-proton interaction but accounting for the interaction between the η -meson and the proton. Due to the lower strength of this interaction the expected deviations from the uniform distributions are by about two orders of magnitude smaller, but one recognizes a slight enhancement of the density in the range of low invariant masses of proton- η subsystems. However, due to weak variations of the proton- η scattering amplitude the enhancement originating from the η -meson interaction with one proton is not separated from the η -meson interaction with the second proton. Therefore an overlapping of broad structures occurs. It is observed that the occupation density grows slowly with increasing s_{pp} opposite to the effects caused by the S-wave proton-proton interaction, yet similar to the modifications expected for the P-wave one [41]. From the above example it is obvious that only in high statistics experiments signals from the meson-nucleon interaction can appear over the overwhelming nucleon-nucleon final state interaction. It is worth noting, however, that the Dalitz plot does not reflect any possible correlations between the entrance and exit channel [39].

The Dalitz plot representation allows also for a simple interpretation of the kinematically available phase-space volume as an area of that plot. Namely, equation (5) becomes:

$$\sigma = \frac{1}{F} \frac{\pi^2}{4s} \int_{(m_1+m_2)^2}^{(\sqrt{s}-m_3)^2} ds_{12} \int_{s_{23}^{min}(s_{12})}^{s_{23}^{max}(s_{12})} ds_{23} |M_{ab \rightarrow 123}|^2, \quad (19)$$

where the limits of integrations defining the boundaries of the Dalitz plot can be expressed as [36]:

$$\begin{aligned} s_{23}^{max}(s_{12}) &= m_2^2 + m_3^2 - \frac{1}{2s_{12}} \left\{ (s_{12} - s + m_3^2)(s_{12} + m_2^2 - m_1^2) - \sqrt{\lambda(s_{12}, s, m_3^2)\lambda(s_{12}, m_2^2, m_1^2)} \right\}, \\ s_{23}^{min}(s_{12}) &= m_2^2 + m_3^2 - \frac{1}{2s_{12}} \left\{ (s_{12} - s + m_3^2)(s_{12} + m_2^2 - m_1^2) + \sqrt{\lambda(s_{12}, s, m_3^2)\lambda(s_{12}, m_2^2, m_1^2)} \right\}. \end{aligned} \quad (20)$$

Thus, the phase-space volume kinematically available for the three-body final state can be written by means of only one integral:

$$V_{ps} = \int dV_{ps} = \frac{\pi^2}{4s} \int_{(m_1+m_2)^2}^{(\sqrt{s}-m_3)^2} ds_{12} \int_{s_{23}^{min}(s_{12})}^{s_{23}^{max}(s_{12})} ds_{23} = \frac{\pi^2}{4s} \int_{(m_1+m_2)^2}^{(\sqrt{s}-m_3)^2} \frac{ds_{12}}{s_{12}} \sqrt{\lambda(s_{12}, s, m_3^2)\lambda(s_{12}, m_2^2, m_1^2)}, \quad (21)$$

whose solution leads, in general, to elliptic functions [36]. However, in the nonrelativistic approximation it has the following closed form:

$$V_{ps} = \frac{\pi^3}{2} \frac{\sqrt{m_1 m_2 m_3}}{(m_1 + m_2 + m_3)^{\frac{3}{2}}} Q^2, \quad (22)$$

where the substitution of the non-relativistic relation between η_{m_3} and Q

$$Q = \frac{m_3^2 + 2 m_3^2 (m_1 + m_2)}{2 (m_1 + m_2)} \eta_{m_3}^2$$

gives the Ss partial cross section of equation (16). On the basis of the formula (22) the kinematically available phase-space volume (V_{ps}) can be as easily calculated as the excess energy Q . Close-to-threshold — in the range of a few tens of MeV — the non-relativistic approximation differs only by a few per cent from the full solution given in equation (21), which in fact with an up-to-date computer can be solved numerically with little effort. Therefore, in the following chapters we will describe the data as a function of V_{ps} as well as of Q or η_M , if it is found to be appropriate.

The choice of the proper observables for picking up the dynamical effects in the case of a four-particle exit channel is by far more complicated. It was for example discussed by Chodrow [42] who has given the extension of the Dalitz representation to the four-body final states. He has proven that the two-body correlation in the case where there are two identical particles in the four-body final state — say ij — can be directly visible in the density distribution of events on the $F_j E_i$ -plane, where E_i denotes the energy of particle i and F_j is defined as:

$$F_j = \frac{1}{2} \left\{ E_j \sqrt{(E_j^2 - m_j^2)} - m_j^2 \cosh^{-1} \left(\frac{E_j}{m_j} \right) \right\}. \quad (23)$$

2.2 Scattering inside the deuteron

Extending meson production experiments from the free nucleon-nucleon interaction to interactions of light nuclei is interesting for many reasons. First of all it offers the possibility to study more complex production processes with more than only one target nucleon involved. Furthermore, in the absence of free neutron targets, it gives the possibility to study the meson creation in the proton-neutron interaction at proton-beam facilities, complementary to a neutron-induced π meson production with use of high-resolution (energy spread less than 1 MeV) secondary neutron beams [43–45] investigated at TRIUMF [46]. The technique based on quasi-free proton-nuclear reactions, with a beam of protons reacting with a nucleus in the target is utilized at the CELSIUS and COSY accelerators [47–51]. First experiments of π^0 meson production in the proton-neutron reaction with simultaneous tagging of the spectator proton have recently been carried out at CELSIUS and resulted in a resolution of the excess energy equal to $\sigma = 1.8$ MeV [47]. In the quasi-free approximation (fig. 5) it is assumed that the bombarding proton interacts exclusively with one nucleon in the target nucleus and that the other nucleons affect the reaction by providing a momentum distribution to the struck constituent only. This assumption is justified if the kinetic energy of a projectile is large compared to the binding energy of the hit nucleus. In fact, as noticed by Slobodrian [52], also the scattering of protons on a hydrogen target, where the protons are bound by molecular forces, may serve as an extreme example of the quasi-free reaction. In that case, although the hydrogen atoms rotate or vibrate in the molecule, their velocities and binding forces are totally negligible with respect to the velocity and nuclear forces operating the scattering of the relativistic proton [52].




Figure 5: Spectator model for a particle production reaction via $pd \rightarrow p_{sp}X$.




Figure 6: (a) Momentum and (b) kinetic energy distribution of the nucleons in the deuteron, generated according to an analytical parametrization of the deuteron wave function [53,54] calculated from the Paris potential [55].

The deuteron is also relatively weakly bound with a binding energy of $E_B \approx 2.2 \text{ MeV}$, which is by far smaller — more than two orders of magnitude in the case of pion and already more than three orders of magnitude in the case of ϕ meson production — than the kinetic energy of the bombarding protons needed for the creation of mesons in the proton–neutron interaction. However, it has to be considered that even the low binding energy of $E_B \approx 2.2 \text{ MeV}$ results in large Fermi momenta of the nucleons which can not be neglected. The momentum and kinetic energy spectra of the nucleons in the deuteron are shown in figure 6.

In the considered approximation the Fermi motion of the nucleons is the only influence of the internucleon forces on the proton–neutron reaction and hence the struck neutron is treated as a free particle

$$p \begin{pmatrix} n \\ p \end{pmatrix} \longrightarrow p \begin{matrix} n \text{ Meson} \\ p_{sp} \end{matrix} \quad (24)$$

in the sense that the matrix element for quasi-free meson production off a bound neutron is identical to that for the free $pn \rightarrow pn \text{ Meson}$ reaction. The proton from the deuteron is considered as a spectator which does not interact with the bombarding particle, but rather escapes untouched and hits the detectors carrying the Fermi momentum possessed at the moment of the collision. From the measurement of the momentum vector of the spectator proton $\vec{p}_{sp}$ one can infer the momentum vector of the neutron $\vec{p}_n = -\vec{p}_{sp}$ at the time of the reaction and hence calculate the excess energy Q for each event, provided that the beam momentum is known. As an example, a distribution of the excess energy in a quasi-free $pn \rightarrow pn\eta'$ reaction is presented in figure 7a [51]. Due to the large centre-of-mass velocity ($\beta \approx 0.75$) with respect to the colliding nucleons, a few MeV wide spectrum of the neutron kinetic energy inside a deuteron (fig. 6b) is spread by more than a factor of 30.




Figure 7: (a) Distribution of the excess energy Q for the $pn\eta'$ system originating from the reaction $pd \rightarrow p_{sp}pn\eta'$ calculated with a proton beam momentum of 3.350 GeV/c and the neutron momentum smeared out according to the Fermi distribution shown in figure 6a. (b) Spectrum of the off-shell mass of the interacting neutron, as calculated under the assumption of the impulse approximation.

Therefore, especially in the case of near-threshold measurements, where the cross section grows rapidly with increasing excess energy (see e.g. fig. 2) the total centre-of-mass energy $\sqrt{s}$ has to be determined on an event-by-event level. For this purpose, in experiments, the spectator protons are usually registered by silicon pad- or μ -strip detectors [47, 51, 56] which determine their kinetic energy (T_{sp}) and polar emission angle (θ). Thus, it is useful to express s as function of these variables:

$$s = |\mathbb{P}_p + \mathbb{P}_n|^2 = s_0 - 2 T_{sp} (m_d + E_p) + 2 p_p \sqrt{T_{sp}^2 + 2 m_p T_{sp}} \cos(\theta) \quad (25)$$

with s_0 denoting the centre-of-mass energy, assuming a vanishing Fermi motion. Measuring both the energy and the emission angle of the spectator protons it is possible to study the energy dependence of a meson production cross section from data taken at only one fixed beam momentum.

It must be noted, however, that in the framework of the impulse approximation, illustrated in figure 5, the measured spectator proton is a physical particle, yet the reacting neutron is off its mass shell, where the explicit expression for its four-momentum vector $\mathbb{P}_n$, in the rest frame of the deuteron, reads:

$$\mathbb{P}_n = (m_d - m_p - T_{sp}, -\vec{p}_{sp}), \quad (26)$$

with T_{sp} and $\vec{p}_{sp}$ denoting the kinetic energy and the momentum vector of a spectator proton, respectively. The mass spectrum of the interacting neutron ($m_n^2 = |\mathbb{P}_n|^2$) resulting from the distribution of Fermi momentum is shown in figure 7b. It can be seen that the maximum of this spectrum differs only by about $3 \text{ MeV}/c^2$ from the free neutron mass ($m_n = 939.57 \text{ MeV}/c^2$), however on the average it is off by about $9 \text{ MeV}/c^2$. In the frame of the discussed approximation, the struck neutron is never on its mass shell and the minimum deviation from the real mass occurs for vanishing Fermi-momentum and — as can be inferred from equation (26) — is equal to the binding energy $E_B = m_d - m_n - m_p$. Measurements performed at the CELSIUS and TRIUMF accelerators for the $pp \rightarrow pp\eta$ [50] and $pp \rightarrow d\pi^+$ [57] reactions, respectively, have shown that within the statistical errors there is no difference between the total cross section of the free and quasi-free processes. In figure 8a the production of the η meson in free proton-proton collisions is compared to the production inside a deuteron and in the overlapping regions the data agree within the statistical errors. These observations allow to anticipate that indeed the assumption of the identity for the transition matrix element for the meson production off free and quasi-free nucleons bound in the deuteron is correct. In case of the meson production off the deuteron, one can also geometrically justify the assumption of the quasi-free scattering since the average distance between the proton and the neutron is in the order of ⁶ 3 fm. Of course, the other nucleon may scatter the incoming proton and the outgoing meson. However, these nuclear phenomena are rather of minor importance in case of the production on the neutron bound in the deuteron, but should be taken into account for derivations of total cross sections from experimental data.

The reduction of the beam flux on a neutron, due to the presence of the proton, referred to as a shadow effect, decreases for example the total cross section by about 4.5 % [59] for the η -meson production. Similarly, the reduction of the total cross section due to the reabsorption of the outgoing η meson on the spectator proton was found to be only about 3 % [59]. The appraisals were performed according to a formula derived in reference [60] which shows that the cross section for the deuteron reduces by a factor of:

$$R = 1 - \sigma_{\eta N}^{inel} \langle r^{-2} \rangle / 4\pi \quad (27)$$

compared to the free nucleon cross sections. Here $\sigma_{\eta N}^{inel}$ denotes the ηN inelastic cross section and $\langle r^{-2} \rangle$ stands for the average of the inverse square nucleon separation in the deuteron taking the nucleon size into account [59]. The latter effect for the production of mesons like $(\pi, \omega, \eta', \phi)$ is expected to be much smaller, since (as shall be discussed in section 3) the s-wave interaction of the η -meson with nucleons is by far stronger than for any of the mentioned ones.

⁶The matter radius of the deuteron amounts to $\approx 2 \text{ fm}$ [58].




Figure 8: (a) Total cross sections for the $pp \rightarrow pp\eta$ reaction as a function of excess energy measured for the scattering of protons in vacuum (open symbols) [61–66] and inside a deuteron (filled symbols) [50]. (b) Close-to-threshold total cross sections for the reactions $pn \rightarrow d\pi^0$ (squares [44], diamonds [67]) and $pd \rightarrow pd\pi^0$ (circles [68], triangles [69]). The solid curve illustrates the predictions for the total cross section of the $pd \rightarrow pd\pi^0$ reaction assuming that the process is utterly due to the quasi-free $pn \rightarrow d\pi^0$ reaction and taking into account the final state interaction between proton and deuteron. Dashed lines shows the result when the final state interaction was neglected. In the calculations the cross section of the $pn \rightarrow d\pi^0$ was parametrized as presented by the dotted line.

Due to the relatively large distance between the nucleons in the deuteron one could argue that the quasi-free reactions will dominate the production of mesons even with a deuteron in the final state. The conviction that the production of mesons in nucleon–deuteron collisions proceeds predominantly via an elementary scattering of nucleons is also based on the comparative investigations of the π^0 –meson production in proton–neutron and proton–deuteron collisions [70]. The total cross section for the $pd \rightarrow pd\pi^0$ reaction close to the kinematical threshold is by more than two orders of magnitude lower compared to the elementary $pn \rightarrow d\pi^0$ process (see figure 8b). This observation has been successfully explained assuming that the production of the π^0 meson in the $pd \rightarrow pd\pi^0$ reaction is proceeding entirely due to the quasi-free $pn \rightarrow d\pi^0$ reaction [70] for η_M larger than 0.7 (dashed line). In the threshold region the result of the calculation (solid line) describes exactly the experimental data when the proton–deuteron FSI is additionally taken into account. The large difference in magnitude between the threshold cross section for the $pn \rightarrow d\pi^0$ and the $pd \rightarrow pd\pi^0$ reaction can be explained since the incident proton momentum needed to create a π^0 meson via the $pd \rightarrow pd\pi^0$ process is much below the kinematical threshold for the production in elementary proton–neutron collisions. Therefore only nucleons possessing large Fermi momenta can contribute to the π^0 production and this appears with a small probability (compare fig. 6a). The other possible elementary processes $pp \rightarrow pp\pi^0$ and $pn \rightarrow pn\pi^0$ can be neglected since they are expected to lead predominantly to a four-body exit channel. This explanation could serve as a heuristic example, however recent data [69] on the cross sections of $dp \rightarrow dp\pi^0$ show that the supposed spectator protons to the elementary $np \rightarrow d\pi^0$ reaction are observed in phase-space regions corresponding to zero Fermi momenta and hence the quasi-free production could be significant only at energies close to the $NN \rightarrow NN\pi$ threshold ($\eta_M(pd \rightarrow pd\pi^0) \approx 0.83$). In this experiment the deuteron beam was impinging onto the hydrogen target, and hence the spectator protons were boosted in the laboratory permitting registration also of the low-energy part of the spectators’ spectral distribution.

A natural extension of the meson production in quasi-free proton–neutron collisions with either a deuteron used as the target or as the beam is the combination of both methods into a double quasi-free production occurring during the collisions of deuterons as depicted in figure 9.




Figure 9: Schematic depiction of the double quasi-free $nn \rightarrow nnX$ reaction. During the collision of deuterons (left hand side of the figure, with the total momentum (solid arrow) resulting from the sum of the beam momentum (dotted arrow) plus the Fermi momentum (short arrow)) a double quasi-free neutron–neutron reaction may lead to the creation of mesons (small gray circle). The spectator protons (black circles) leave the reaction region with their initial momentum plus the Fermi momentum, which they possessed at the moment of the reaction. Neutrons are plotted as open circles. Due to the large relative momenta between spectators and the outgoing neutrons ($\sim 1 \text{ GeV}/c$ close to the threshold for the η meson production) a distortion of the nnX system by the accompanying protons can be neglected.

Utilizing this technique [71] and registering both the slow and fast spectator protons could allow for the study of meson production in neutron–neutron collisions with a very precise determination of the excess energy which depends on the accuracy for the registration of the momentum vectors of the spectators.

3 Phenomenology of the initial and final state interaction

The simplest method to study the interactions between meson and nucleon is to scatter a beam of mesons off hydrogen or deuteron targets and to measure the behaviour of the total and differential cross section in the elastic scattering of the considered particles. Figure 10a presents the total cross section for K^+p and K^-p elastic scattering as a function of the centre-of-mass excess energy. When comparing the data to calculations [72] — including the changes of phase-space integral (eq. (10)) and Coulomb interaction in the initial and final states — one observes a huge enhancement for the K^-p cross section with decreasing excess energy and a slight suppression in the case of K^+p scattering. This observation may be attributed to the slight repulsion due to the kaon–proton hadronic interaction and the significantly larger scattering caused by the strong interaction between the K^- and proton due to the vicinity of the $\Lambda(1405)$ hyperon resonance. The effect has a direct influence on the effective mass splitting of kaons immersed in dense nuclear matter [73, 74].




Figure 10: (a) Cross section for the K^+ -proton [75] and K^- -proton [76] elastic scattering. Solid and dashed lines represent the changes of the phase-space integral (eq. (10)) modified by the initial and final state Coulomb interaction ($\sigma_{Kp \rightarrow Kp} = const \cdot C^4 / (16\pi s)$). Both curves are normalized to points of large excess energies. (b) Total cross section of the $pp \rightarrow pp K^+ K^-$ reaction [77, 78]. Statistical and systematical errors are separated by dashes. The solid line indicating calculations of reference [79] is described in the text. (c) Decomposition of the total cross section of the $pp \rightarrow pp \pi^0$ reaction into Ss, Ps and Pp final state angular momenta. The dashed and solid lines represent the η_M^6 and η_M^8 dependence of Ps and Pp partial cross sections, respectively. The remainder is indicated as the dotted line. The Sp partial wave is forbidden by the conservation laws and the Pauli excluding principle. Note that at $\eta_M = 1$ the Pp and Ps partial waves seem to dominate. However, the analysis of the differential cross section measured at CELSIUS at $\eta_M = 0.449$ [80, 81] showed that also a d-wave pion production — due to the interference between Ss and Sd states — constitutes 7% of the total cross section, when a meson-exchange model is assumed. The figure has been adapted from [82].

A Coulomb repulsion was taken into account by multiplying the expression (10) by the Coulomb penetration factor C^2 for both exit and entrance channels. C^2 determines the ratio of the probability of finding two particles close together to the probability of finding two uncharged particles together, all other things being equal [83] and can be expressed as [84]:

$$C^2 = \frac{2\pi\eta_c}{e^{2\pi\eta_c} - 1}, \quad (28)$$

where η_c is the relativistic Coulomb parameter, which for the collision of particles i, j reads:

$$\eta_c = \frac{q_i q_j \alpha}{v} = q_i q_j \alpha \frac{s_{ij} - m_i^2 - m_j^2}{\sqrt{\lambda(s_{ij}, m_i^2, m_j^2)}},$$

with the fine structure constant α , the relative velocity v of the colliding particles and with q_i, q_j denoting their charges ⁷. The comparison of the K^-p scattering data with the phase space integral presented in figure 10a aims only to illustrate the qualitative influence of the meson-nucleon dynamics on the total cross section energy dependence. It is worth noting, however, that at present the data can be qualitatively well described in the frame of the effective chiral Lagrangian with a coupled channel potential [26, 87, 88].

⁷For collisions at an angular momentum l larger than $0\hbar$ the C^2 of equation (28) needs to be multiplied by a factor of $\prod_{n=1}^l (1 + (\eta_c/n)^2)$ [85, 86].

Another method permitting such investigations is the production of a meson in the nucleon–nucleon interaction close to the kinematical threshold where the outgoing particles possess small relative velocities. The experimental observables are the excitation functions and the double differential cross section. A possible deviation from the expectation based on the assumption of homogeneous phase-space abundance corrected for the known nucleon–nucleon interaction delivers information about the meson–nucleon forces. For example, figure 10b presents the first data points on the close-to-threshold production of the K^- meson via the $pp \rightarrow ppK^+K^-$ reaction. The superimposed solid line depicts the results of calculations [79] taking into account the changes of the production amplitude as deduced from the K^+p and K^-p elastic scattering shown in figure 10a, but neglecting the influence of the dominant proton–proton interaction! On the other hand the proton–proton FSI can not be neglected in case of three-body final states, e.g. $pp \rightarrow pp\pi^0$ [8, 89], $pp \rightarrow pp\eta$ [61, 63], or $pp \rightarrow pp\eta'$ [37, 90], since it has strong influence on the total cross section energy dependence enhancing it by more than an order of magnitude for excess energies below $Q \approx 15$ MeV as shall be shown below. Thus it is surprising that in spite of its neglect one can describe the data of the $pp \rightarrow ppK^+K^-$ reaction. The origin of that effect will be investigated experimentally in the near future [91]. At present one can only speculate whether it is due to the partial compensation of the pp and K^-p hadronic interaction or maybe due to the additional degree of freedom given by the four-body final state [92].




Figure 11: (a) The 1S_0 and 3P_0 phase-shifts of the nucleon–nucleon potential shown versus the centre-of-mass kinetic energy available in the proton–proton system. The values have been extracted from the SAID data base [93] (solution SM97). For higher energies the S- and P-wave phase-shifts are nearly the same. This is because the collision parameter required to yield the angular momentum of $1\hbar$ diminishes significantly below 1 fm with increasing energy and consequently the interaction of nucleons — objects of about 1 fm size — becomes almost central. (b) The variable η_M as a function of the excess energy for π^0 , η and η' mesons produced via $pp \rightarrow pp \text{ Meson}$ reactions. (c) The phase-space volume V_{ps} (defined by equation (21)) versus the excess energy Q . The picture indicates that for the production of “heavy mesons” in the nucleon–nucleon interaction at a given Q value there is only a slight difference of the available phase-space volume on the produced meson mass, which is larger than that of π^0 production by about 30 % only. Therefore, for the comparative studies of the production dynamics of different mesons, Q is as much a suitable variable as V_{ps} .

In order to demonstrate the influence of the nucleon–nucleon interaction on the cross section dependence on the excess energy let us consider the production of neutral pseudoscalar mesons (π^0 , η , η') in the collision of protons. Due to the short life-time of these mesons the study of their low-energy interaction with nucleons in direct scattering experiments is not feasible. In general for a three-body exit channel one expects a dependence of the total cross section on the energy which can be described

by the linear combination of partial cross sections of equation (16). Therefore, to be able to extract the information about the final state interaction of the outgoing particles from the energy dependence it is necessary to know precisely the contribution originating from different partial waves. Yet appropriately close-to-threshold there is only one important combination of the angular momenta of emitted particles (Ss) and in this region the energy dependence of the total cross section is uniquely determined. However, the range of excess energies for the S-wave dominance changes strongly with the mass of the produced meson.

For the production of the π^0 meson in proton-proton collisions the investigations with polarized beam and targets [82,94] allowed to deduce that the Ss partial-wave accounts for more than 95 % of the total cross section up to $\eta_M \approx 0.4$, as can be seen in figure 10c, where the Ss contribution is indicated by the dotted line. The Ss contribution was inferred assuming the η_M^6 and η_M^8 dependence for Ps and Pp partial waves, respectively. These are power-laws taken from proportionality (16), which was derived under the assumption of non-interacting particles. Relatively small values of 3P_0 -wave nucleon-nucleon phase-shifts at low energies (compared to 1S_0 phase-shifts in figure 11a) and similarly weak low-energy interactions of P-wave protons in other spin combinations [95] justify this assumption.

The measurements with the polarized beam and target allow a model independent determination of the contribution from individual partial waves. In case of the $\vec{p}\vec{p} \rightarrow pp\pi^0$ reaction, due to the identical particles in the initial state and the rotational equivalences [82,96] there are only seven independent polarization observables, e.g. two beam analyzing powers and five linear combination of spin correlation coefficients.

In particular, as shown in references [82,94] the close to threshold contributions of the Ps and Pp partial waves can be determined in the model-free way from the measurements of the spin dependent total cross sections only. For example the strength of the Ps final state can be expressed as [82]:

$$\sigma(Ps) = \frac{1}{4} \left(\sigma_{tot} + \Delta\sigma_T + \frac{1}{2}\Delta\sigma_L, \right) \quad (29)$$

where σ_{tot} denotes the total unpolarized cross section and $\Delta\sigma_T$ and $\Delta\sigma_L$ stand for differences between the total cross sections measured with anti-parallel and parallel beam and target polarizations. Subscripts T and L associate the measurements with the transverse and longitudinal polarizations, respectively.

In accordance with the phenomenology of Gell-Mann and Watson [4] described in the previous section one expects that also in the case of heavier mesons the Ss partial wave combination will constitute the overwhelming fraction of the total production cross section for η_M smaller than 0.4. This implies — as can be deduced from the relation between η_M and Q illustrated in figure 11b — that mesons heavier than the pion are produced exclusively via the Ss state in a much larger excess energy range and hence larger phase-space volume (see figure 11c). In particular in case of η' this is larger by more than one order of magnitude. Thus, whereas in case of π^0 the onset of higher partial waves is observed at Q around 10 MeV it is expected only above 100 MeV and above ≈ 40 MeV for η' and η mesons, respectively. Figures 12b and 12c present the angular distributions of the created η meson in proton collisions. It is evident that at $Q = 15.5$ MeV and still at $Q = 41$ MeV the production of the η meson is completely isotropic within the shown statistical errors. Although at $Q = 41$ MeV the accuracy of the data does not exclude a few per cent of the contribution originating from higher partial waves, the dominance of the s-wave creation is evident. Similarly, the measurements of the differential cross section (fig. 12a) for the $pp \rightarrow pp\eta'$ reaction performed at SATURNE [97] at $Q = 143.8$ MeV are still consistent with pure Ss-wave production, though the relatively large error bars would allow for other contributions on a few per cent level ($\approx 10\%$).




Figure 12: Differential cross section of the $pp \rightarrow pp \text{Meson}$ reaction as a function of the meson centre-of-mass polar angle. Dashed lines indicate the isotropic distribution. Shown are results of measurements for the $pp \rightarrow pp\eta'$ reaction taken at $Q = 143.8 \text{ MeV}$ [97] (a) and for the $pp \rightarrow pp\eta$ reaction at $Q = 15.5 \text{ MeV}$ [66] (b) and $Q = 41 \text{ MeV}$ [98] (c). In all pictures only statistical errors are plotted, which in figure (b) are smaller than the size of symbols. The distribution presented in picture (b) is consistent with a measurement performed at an excess energy of $Q = 16 \text{ MeV}$ by means of the PROMICE/WASA detector [99], whereas the data at $Q = 37 \text{ MeV}$ also from reference [99] deviate significantly from isotropy. However, data shown in picture (c) have been taken with a detector of much higher angular acceptance.

Table 2: Momentum transfer Δp calculated according to equation 17 and the corresponding distance $R \approx \hbar/\Delta p$ probed by the $NN \rightarrow NN \text{Boson}$ reaction at the kinematical threshold for different particles produced. The table has been adapted from [20].

particle	mass [MeV]	$\Delta p [\text{fm}^{-1}]$	$R [\text{fm}]$
γ	0	0.0	∞
π	140	1.9	0.53
η	550	3.9	0.26
ρ, ω	780	4.8	0.21
η'	960	5.4	0.19
ϕ	1020	5.6	0.18

Let us now consider to what extent the energy dependence of the total cross section in the estimated range of the dominance of the Ss partial waves can be understood in terms of the phase-space variation and the interaction between the particles participating in the reaction. Watson [18] and Migdal [100] have proposed the factorization of the amplitude in the case where the production is of short range and the interaction among the outgoing particles is of long range. This requirement is well fulfilled in the case of the close-to-threshold meson production due to the large momentum transfer (Δp) between the interacting nucleons needed to create the considered mesons ($\pi, \eta, \dots, \phi$). According to the Heisenberg uncertainty relation the large momentum transfer brings about a small space in which the primary creation of the meson takes place. In table 2 the distance probed by the $NN \rightarrow NN \text{Meson}$ reaction at threshold is listed for particular mesons. It ranges from 0.53 fm for pion production to 0.18 fm for the ϕ meson, whereas the typical range of the strong nucleon-nucleon interaction at low energies determined by the pion exchange may exceed a distance of few Fermi and hence is by one order of magnitude larger

than the values listed in table 2. Thus in analogy to the Watson–Migdal approximation for two–body processes [18] the complete transition matrix element of equation (5) may be factorized approximately as ⁸

$$|M_{pp \rightarrow ppX}|^2 \approx |M_{FSI}|^2 \cdot |M_0|^2 \cdot F_{ISI}, \quad (30)$$

where M_0 represents the total short range production amplitude, M_{FSI} describes the elastic interaction among particles in the exit channel and F_{ISI} denotes the reduction factor accounting for the interaction of the colliding protons. Further, in the first order approximation one assumes that the particles are produced on their mass shell and that the created meson does not interact with nucleons. This assumption implies that the $|M_{FSI}|^2$ term can be substituted by the square of the on–shell amplitude of the nucleon–nucleon elastic scattering:

$$|M_{FSI}|^2 = |M_{NN \rightarrow NN}|^2. \quad (31)$$

Effects of this rather strong assumption will be considered later, while comparing the estimation with the experimental data.

In the frame of the optical potential model the scattering amplitude is determined by phase–shifts. Particularly, the 1S_0 proton–proton partial wave — relevant for further consideration — can be expressed explicitly as follows [101]:

$$M_{pp \rightarrow pp} = \frac{e^{-i\delta_{pp}(^1S_0)} \cdot \sin \delta_{pp}(^1S_0)}{C \cdot k}, \quad (32)$$

where C denotes the square root of the Coulomb penetration factor defined by equation (28), k stands for either proton momentum in the proton–proton rest frame and the phase–shift is indicated by δ_{pp} . The phase–shifts $\delta_{pp}(^1S_0)$ can be extracted from the SAID data base (see fig. 11a) or, alternatively, can be calculated according to the modified Cini–Fubini–Stanghellini formula including the Wong–Noyes Coulomb correction [102–104]

$$C^2 k \operatorname{ctg}(\delta_{pp}) + 2k\eta_c h(\eta_c) = -\frac{1}{a_{pp}} + \frac{b_{pp}k^2}{2} - \frac{P_{pp}k^4}{1 + Q_{pp}k^2}, \quad (33)$$

where $h(\eta_c) = -\ln(\eta_c) - 0.57721 + \eta_c^2 \sum_{n=1}^{\infty} \frac{1}{n \cdot (n^2 + \eta_c^2)}$ [83], with η_c the same as in eq. (28).

The phenomenological quantities $a_{pp} = -7.83$ fm and $b_{pp} = 2.8$ fm denote the scattering length and effective range [102], respectively. The parameters $P_{pp} = 0.73$ fm³ and $Q_{pp} = 3.35$ fm² are related to the detailed shape of the nuclear potential and derived from a one–pion–exchange model [102]. Substituting equation (33) into equation (32) allows to calculate the low–energy amplitude for the proton–proton elastic scattering ⁹:

$$|M_{pp \rightarrow pp}|^2 = \frac{C^2}{C^4 k^2 + \left(-\frac{1}{a_{pp}} + \frac{b_{pp}k^2}{2} - \frac{P_{pp}k^4}{1 + Q_{pp}k^2} - 2k\eta_c h(\eta_c) \right)^2}. \quad (34)$$

The result is presented as a solid line in figure 13a and is in good agreement with the values obtained from the phase–shifts of the VPI partial wave analysis [93], shown as solid circles and with the phase–shifts of the Nijmegen analysis [105], shown as open squares.

⁸For a comprehensive discussion of the FSI and ISI issue including a historical overview and a criticism of various approaches the reader is referred to [22].

⁹In principle the formula is valid for $k \leq 133$ MeV/c [103]. $|M_{pp \rightarrow pp}|^2$ is taken to be constant for larger values of k .




Figure 13: (a) Square of the proton–proton scattering amplitude versus k , the proton momentum in the proton–proton subsystem from [101, 102] (solid line), [106] (dashed line), and [107, 108] (dotted line). The filled circles are extracted from [93] and the opened squares from [105]. The curves and symbols are arbitrarily normalized to be equal at maximum to the result from reference [106], shown as the dashed line. (b) Total cross section for the $pp \rightarrow pp \pi^0$ reaction as a function of the centre–of–mass excess energy Q . Data are from refs. [8, 81, 89, 109]. The dashed line indicates a phase–space integral normalized arbitrarily. The phase–space distribution with inclusion of proton–proton strong and Coulomb interactions fitted to the data at low excess energies is shown as the solid line. The dotted line indicates the parametrization of reference [110] written explicitly in equation (35), with $\epsilon = 0.3$.

The factor C^2 is always less than unity due to the Coulomb repulsion between protons. At higher energies, where C^2 is close to unity, the nuclear scattering will be predominant and for the very low energies the Coulomb and nuclear interactions are competing. The Coulomb scattering dominates approximately up to about 0.8 MeV of the proton energy in the rest frame of the other proton, where C^2 equals to one–half [83]. In the case of the $pp \rightarrow pp \text{ Meson}$ reaction the maximum possible energy of a proton seen from another proton is equal to 0.8 MeV already at an excess energy of about $Q = 0.4$ MeV.




Figure 14: (a) Total cross section for the $pp \rightarrow pp \eta'$ reaction as a function of the centre–of–mass excess energy Q . Data are from refs. [62, 90, 97, 111, 112]. Statistical and systematical errors are separated by dashes. The solid line shows the phase–space distribution with inclusion of proton–proton strong and Coulomb interactions. The dotted line indicates the parametrization of reference [110] written explicitly in equation (35), with $\epsilon = 0.3$ and the dashed line indicates a phase–space integral normalized arbitrarily. (b) Stars represent the data of the $pp \rightarrow p K^+ \Lambda$ reaction [113–119]. The dashed line presents the energy dependence defined by the phase–space. The solid line shows the calculation with proton– K^+ Coulomb repulsion and a proton– Λ strong interaction taken into account as described in the text.

Therefore, a significant influence of the Coulomb repulsion on the energy dependence of the total production cross section is expected only at very low excess energies, i.e. conservatively for $Q \leq 2 \text{ MeV}$.

Assuming that the on-shell proton-proton amplitude exclusively determines the phase-space population one can obtain the total cross section energy dependence substituting equation (34) into the formula (19). Solid lines in figures 13b and 14a represent the determined dependence for the $pp \rightarrow pp\pi^0$ and $pp \rightarrow pp\eta'$ reactions, respectively. The absolute scale of the calculations has been fixed by the normalization to the data. One recognizes the good agreement with the experimental points in the excess energy range up to $Q \approx 10 \text{ MeV}$ for π^0 meson production and up to $Q \approx 150 \text{ MeV}$ for η' . This confirms the earlier considerations that the domain of Ss partial wave dominance should widen significantly with the mass of the created meson and as expected it extends above 100 MeV in case of the η' meson. Comparing the data to the arbitrarily normalized phase-space integral of equation (21) reveals that the proton-proton FSI enhanced the total cross section by more than an order of magnitude for low excess energies. This is particularly clear in figure 14a where the logarithmic scale is utilized. However, the much weaker nucleon-hyperon interaction increases the total cross section for the close-to-threshold K^+ meson production only by about a factor of two. Again in that case — as shown in figure 14b — the factorization of the matrix element into constant on-shell primary production amplitude and the elastic scattering between outgoing baryons leads to the good description of the close-to-threshold data. In this case the Coulomb interaction between K^+ meson and proton was taken into account multiplying the scattering amplitude by the Coulomb penetration factor.

The dotted line in figures 13b and 14a, which for $Q < 50 \text{ MeV}$ is practically indistinguishable from the solid line, presents the excess energy dependence of the total cross section taking into account the proton-proton FSI effects according to the model developed by Fäldt and Wilkin [110,120]. Representing the scattering wave function in terms of bound state wave function the authors derived a closed formula which describes the effects of the nucleon-nucleon FSI as a function of the excess energy Q only. This approach is specifically useful for the description of the spin-triplet proton-neutron FSI, due to the existence of a bound state (deuteron) with the same quantum numbers. Though the bound state of the proton-proton system does not exist, the model allows also to express the total cross section energy dependence for a $pp \rightarrow pp \text{ Meson}$ reaction by a simple and easily utilizable formula:

$$\sigma = \text{const.} \cdot \frac{V_{ps}}{F} \cdot \frac{1}{\left(1 + \sqrt{1 + \frac{Q}{\epsilon}}\right)^2} = \text{const.}' \cdot \frac{Q^2}{\sqrt{\lambda(s, m_p^2, m_p^2)}} \cdot \frac{1}{\left(1 + \sqrt{1 + \frac{Q}{\epsilon}}\right)^2}, \quad (35)$$

where the parameter ϵ has to be settled from the data. The flux factor F and the phase-space volume V_{ps} are given by equations (9) and (22), respectively. The normalization can be determined from the fit of the data which must be performed for each reaction separately.




Figure 15: (a) Total cross section for the $pp \rightarrow pp\eta$ reaction as a function of the centre-of-mass excess energy Q . Data are from refs. [50, 61–66]. Statistical and systematical errors are separated by dashes. The dashed line indicates a phase-space integral normalized arbitrarily. The phase-space distribution with inclusion of proton-proton strong and Coulomb interactions fitted to the data in the excess energy range between 15 and 40 MeV is shown as the solid line. Additional inclusion of the proton- η interaction is indicated by the dotted line. The scattering length $a_{p\eta} = 0.7 \text{ fm} + i 0.4 \text{ fm}$ and the effective range parameter $b_{p\eta} = -1.50 \text{ fm} - i 0.24 \text{ fm}$ [125] have been arbitrarily chosen for the calculations described in the text. (b) Dalitz plot distribution of the $pp \rightarrow pp\eta$ reaction at an excess energy of $Q = 37.6$ MeV, corrected for the detection acceptance. Out of the two invariant masses corresponding to two $p - \eta$ pairs the one being larger is plotted along the x-axis. The figure is taken from [63] (A similar spectrum for the $pp\pi^0$ system [89] reveals the influence of the proton-proton interaction, yet again, as for the total cross section energy dependence (fig. 13b), the proton- π^0 interaction is too weak to affect observably the density distribution of the Dalitz plot.). (c) Dalitz plot distribution of the $pp \rightarrow pp\eta$ reaction at $Q = 15.5$ MeV, corrected for the detection acceptance and the proton-proton FSI [66]. The proton-proton FSI enhancement factor was calculated as square of the on-shell proton-proton scattering amplitude as written in equation (34).

In the case of η meson production the interaction between nucleons is evidently not sufficient to describe the increase of the total cross section for very low and very high excess energies, as can be seen in figure 15a. The solid line in the figure was normalized to the data at an excess energy range between 15 MeV and 40 MeV. The increase of the total cross section for higher energies can be assigned to the outset of higher partial waves. As expected from the previous considerations, this is indeed seen at $Q \approx 40$ MeV where the energy dependence of the total cross section starts to change its shape. On the contrary, the close-to-threshold enhancement — being by about factor of two larger than in the case of the π^0 and η' mesons — can be assigned neither to the contribution from other than Ss partial waves nor to the variation of the primary production amplitude $|M_0|$. The latter is expected to change at most by a few per cent for excess energies below 20 MeV [121]. Instead, this discrepancy can be plausibly explained by the influence of the attractive interaction between the η meson and the proton. Note, that the real part of the scattering length of the η -proton potential — depending on the analysis method and the studied reaction — is 3 to 10 times [122, 123] larger than the scattering length for π^0 -proton scattering ($a_{p\pi} \approx 0.13 \text{ fm}$) [124]. Hence, the modifications of the total cross section energy dependence due to π^0 -proton interaction are too weak to be observed within the up-to-date accuracy of measurements and calculations. However, the influence of the η -proton interaction presented in figure 15 is evident and hereafter it will be considered whether it may serve for the estimation of the η -proton scattering parameters.

The strict quantitative calculation requires, however, the evaluation of the three-body Faddeev

equation ¹⁰. Here, we will rather present a simple phenomenological treatment which shall lead to the qualitative understanding how the mutual interaction among three outgoing particles affects the total cross section dependence on excess energy. One of the simplest possibilities based on the naive probabilistic interpretation of the incoherent pairwise interaction would be to factorize the overall enhancement factor into corresponding pair interactions [34, 127]:

$$|M_{FSI}|^2 = |M_{12 \rightarrow 12}|^2 \cdot |M_{13 \rightarrow 13}|^2 \cdot |M_{23 \rightarrow 23}|^2, \quad (36)$$

where $|M_{ij \rightarrow ij}|^2$ denotes the square of the elastic scattering amplitude of particles i and j . The $|M_{pp \rightarrow pp}|^2$ can be evaluated according to the formula (34), which for the s-wave η -proton scattering, after substitution of $C^2 = 1$ and $\eta_c = P_{pp} = 0$, reduces to ¹¹:

$$|M_{p\eta \rightarrow p\eta}|^2 = \left| \frac{1}{\frac{1}{a_{p\eta}} + \frac{b_{p\eta} k_{p\eta}^2}{2} - i k_{p\eta}} \right|^2, \quad (37)$$

where

$$k_{p\eta} = \frac{\sqrt{\lambda(s_{p\eta}, m_\eta^2, m_p^2)}}{2 \sqrt{s_{p\eta}}}$$

denotes the η momentum in the proton- η rest frame. The scattering length $a_{p\eta}$ and effective range $b_{p\eta}$ are complex variables with the imaginary part responsible e.g. for the $p\eta \rightarrow p\pi^0$ conversion.

The factorization of both i) the overall production matrix element (eq. (30)) and ii) the three particle final state interactions (eq. (36)) applied to the formula (19) gives the following expression for the total cross section of the $pp \rightarrow pp\eta$ reaction:

$$\sigma = \frac{F_{ISI} |M_0|^2}{F} \frac{\pi^2}{4s} \int_{(m_p+m_p)^2}^{(\sqrt{s}-m_\eta)^2} ds_{pp} \int_{s_{p2\eta}^{min}(s_{pp})}^{s_{p2\eta}^{max}(s_{pp})} ds_{p2\eta} |M_{pp \rightarrow pp}(s_{pp})|^2 \cdot |M_{p1\eta \rightarrow p1\eta}(s_{p1\eta})|^2 \cdot |M_{p2\eta \rightarrow p2\eta}(s_{p2\eta})|^2, \quad (38)$$

where the protons are distinguished by subscripts. Exploring formulas (34) and (37) gives the results shown as the dotted line in figure 15a. Evidently, the inclusion of the proton- η interaction enhances the total cross section close-to-threshold by about a factor of 1.5 and leads to a better description of the data. The effect of the proton- η interaction is also seen in the experimental distribution of the differential cross sections $d^2\sigma/(ds_{p1\eta} ds_{p2\eta})$ shown in figures 15b and 15c. These distributions originating from kinematically complete measurements comprise the whole experimentally available information about the interactions of the $pp\eta$ -system. In figure 15b one recognizes the increase of the distribution density at regions of small invariant masses of the proton-proton and proton- η subsystems. At this excess energy ($Q = 37.6$ MeV) these regions are quite well separated. However, since this is close to the energy where the advent of higher partial waves is awaited, the possible contribution from the P-wave proton-proton interaction can not be a priori excluded. Specifically, the latter leads to the enhancement at large invariant masses of the proton-proton pair [41] and hence affects the phase-space region where the modification from the proton- η interaction is expected. Figure 15c presents

¹⁰An exact derivation of the Faddeev equation can be found for example in [126].

¹¹Note that the sign of the term $-1/a$ from equation (34) was changed because the imaginary part of the proton- η scattering length is positive [125], as we also adopted here, whereas in the majority of works concerning nucleon-nucleon interaction, the scattering length is negative [31]. We are grateful for this remark to A. Gasparyan.

the high statistics data taken at an excess energy of $Q = 15.5$ MeV, where the assumption of the total dominance of the Ss partial wave is rather save. At this excess energy, corresponding to the small relative momentum range $k_{p\eta}^{max} \approx 105$ MeV/c, the variations of the proton- η scattering amplitude are quite moderate (fig. 18c) and as expected from simulations presented in the previous section, the increase of the phase-space population due to the η -meson interaction with one proton is not separated from the other one. Interestingly, the density growth with increasing invariant mass of the proton-proton system is much faster than expected from the simulations presented in figure 4d, which have been performed with a scattering length equal to $a_{p\eta} = 0.7 \text{ fm} + i 0.3 \text{ fm}$. However, when reducing the proton-proton FSI effect to a multiplicative factor, one finds that it depends on the assumed nucleon-nucleon potential and on the produced meson mass [128].

This issue was recently vigorously investigated e.g. by authors of references [20–22, 128, 129] and we shall briefly report this here as well. Up to now we factorized the transition matrix element into a primary production of particles and its on-shell rescattering in the exit channel (eqs. (30)(31)(36)). Though it is a crude approximation, neglecting the off-shell effects of the production process completely, it astoundingly leads to a good description of the energy dependence of the total cross section, as already demonstrated in figures 13b, 14a,b and 15a. The off-shell effects, as pointed out by Kleefeld [22], could have been safely neglected in case of the electromagnetic transitions in atoms or β decays, where the excitation energy of the involved nucleons is by many orders of magnitude smaller than their masses and the initial and final states go hardly off-shell [22]. However, in case of the $NN \rightarrow NN \text{ Meson}$ process the large excitation energy of the colliding nucleons is comparable with the nucleon masses and the primary interaction may create the particles significantly far from their physical masses, so that the off-shell effects can not be a priori disregarded.




Figure 16: Diagrammatic representation of the DWBA expressed by equation (39). (a) The primary production term. (b) The loop diagram including the nucleon-nucleon FSI. $T(k', k)$ stands for the half-off-shell T matrix with k and k' denoting the on- and off-shell centre-of-mass momentum in the nucleon-nucleon system, respectively. q indicates the momentum of the created meson in the reaction centre-of-mass and p the momentum of the colliding nucleons.

Generally, the decomposition of the total production amplitude into the primary production and the subsequent nucleon-nucleon interaction visualized in figure 16 is expressed by the formula:

$$M = M_0^{on} + M_0^{off} G T_{NN}, \quad (39)$$

where the second term of the equation represents the integration over the intermediate (k') momenta of the off-shell production amplitude and the half-off-shell nucleon-nucleon T matrix [128]. Assuming

that the primary production occurs in such a way that one nucleon emits the meson which then re-scatters on the other nucleon and appears as a real particle the authors of reference [128] found that the enhancement of the cross section due to the nucleon–nucleon interaction depends strongly on the mass of the created meson. This is because with the increasing mass of the produced meson the distance probed by the nucleon–nucleon interaction decreases (see table 2) and hence the relevant range of the off-shell momenta becomes larger. The effect for the $pp \rightarrow pp$ Meson reactions is presented in figure 17a, where one can see that, when utilizing the Bonn potential model for the nucleon–nucleon T matrix, the enhancement in case of the π^0 production is by about a factor of four larger than for the η or η' mesons. A similar conclusion, but with the absolute values larger by about 40 %, was drawn for the Paris NN potential [128]. On the contrary, when applying the Yamaguchi potential into calculations the enhancement grows with the increasing mass of the meson, as shown in figure 17b. Thick solid curves in figures 17a and 17b show the results of the frequently applied approximation of the nucleon–nucleon FSI effects:

$$M = M_0^{on} + M_0^{off} G T_{NN} \approx M_0^{on} \cdot (1 + G T_{NN}) = M_0^{on} J^{-1}(-k) \equiv M_0^{on} F_{NN}(k), \quad (40)$$

where the overall transition matrix element M is factorized to the primary on-shell production and the NN FSI expressed as the inverse of the Jost function $J^{-1}(-k)$ [108]. As can be seen in figures 17a and 17b, the variation of the absolute values — of such obtained enhancement factors — with the applied potential is severe.




Figure 17: The FSI factor for Bonn [131] a) and Yamaguchi [132] b) potentials. The solid, dotted and dashed lines correspond to π^0 , η and η' meson production, respectively. Thick solid lines indicate the inverse of the squared Jost function ($|F_{pp}(k)|^2 = |J(-k)|^{-2}$). c) The FSI factors for π^0 production of Paris (dotted curve) [55], Bonn (dashed curve) and Yamaguchi (solid curve) potentials normalized to be equal at maximum to the inverse of the squared Jost function of the Yamaguchi potential (thick solid line). The shapes stemming from different potentials are almost indistinguishable. Note that the thick solid line corresponds to the dashed line in figure 13a. The figures are adapted from reference [128].

Since the physical value of the total cross section can not depend on the off-shell features of the potential used in calculations, which are in principal not measurable [130], the differences in the magnitude of the $|F_{pp}|$ factor must reflect itself in a corresponding dependence of the primary production amplitude on the potential used. Therefore, it is of crucial importance to realize that the values of the threshold amplitudes $|M_0|$ are significant only in the context of the potential they were extracted from. Figure 17c demonstrates, however, that the shapes of the enhancement factors, with the meson

exchange mechanism assumed for the primary production, are pretty much the same, independently of the applied NN potential and correspond to the form of the Jost function inferred from the Yamaguchi potential. This indicates that the energy dependence of the NN FSI factors is predominantly determined by the on-shell NN T matrix.

In references [20–22] the formula for the transition matrix element with explicit dependence on the considered off-shell features for the initial and final state interaction is derived:

$$M = \left\{ 1 + \frac{1}{2} [\eta(k)e^{2i\delta(k)} - 1] \cdot [1 + P_f(p, k)] \right\} M_0 \left\{ 1 + \frac{1}{2} [\eta(p)e^{2i\delta(p)} - 1] \cdot [1 + P_i(p, k)] \right\}, \quad (41)$$

where subscripts i and f indicate the initial and final state, respectively. The functions $P(p, k)$ exhibit all the off-shell effects of the NN interaction and the primary production current [20] and δ and η denote the phase-shift and inelasticity, correspondingly. At threshold, the inelasticity in the exit channel is equal to unity ($\eta(k) = 1$) due to the small relative momentum of the outgoing nucleons. The last term of the formula expresses the influence of the initial state interaction on the production process. Due to the large relative momenta of the colliding protons needed to create a meson it is characterized by a weak energy dependence in the excess energy range of a few tens of MeV. For example in figure 11a one can see that the phase-shift variation of the 3P_0 partial wave (having predominant contribution to the threshold production of pseudoscalar mesons) in the vicinity of threshold for mesons heavier than π^0 is indeed very weak. Taking additionally into account that the initial state off-shell function $P_i(p, k)$ is small (at least it is the case for meson exchange models [21]) one reduces the influence of the NN initial state interaction to the reduction factor F_{ISI} which can be estimated from the phase-shifts and inelasticities only:

$$F_{ISI} = \frac{1}{4} \left| \eta(p) e^{2i\delta(p)} + 1 \right|^2. \quad (42)$$

At the threshold for π meson production this is close to unity since at this energy the inelasticity is still nearly 1 and the 3P_0 phase-shift is close to zero (see figure 11a). However, at the η threshold, where the phase-shift approaches its minimum, the proton–proton ISI diminishes the total cross section already by a factor of 0.2 [21]. A similar result was obtained using a meson exchange model for η production in the $pp \rightarrow pp\eta$ reaction and calculating the proton–proton distortion from the coupled-channel πNN model [19]. The authors of reference [19] concluded that the initial proton–proton distortion reduces the total cross section by about a factor of ≈ 0.26 , which keeps constant at least in the studied range of 100 MeV in kinetic beam energy. Hence, the closed formula (42) disregarding the off-shell effects ($P_i(p, k)$) permits to estimate the cross section reduction due to the initial state distortion with an accuracy of about 25 %. The shape of the 3P_0 phase-shift shown in figure 11a indicates that the effect is at most pronounced close to the η production threshold, yet for the η' meson the formula (42) leads to a factor $F_{ISI} = 0.33$ [133]. The primary production amplitude as well as the off-shell effects of the nucleon–nucleon FSI ($P_f(p, k)$) are also weakly energy dependent [20], since they account for the short range creation mechanism, which shall be considered in the next section. The accordance of the experimental data with the simple factorization represented by solid lines in figures 13b, 14 and 15a fully confirms the above considerations which imply that the total cross section energy dependence and the occupation density on a Dalitz plot is in the first order determined by the on-shell scattering of the outgoing particles. However, since the distortion caused by the nucleons is by some orders of magnitude larger than that due to the meson–nucleon interaction, even small fractional inaccuracies in the description of nucleon–nucleon effects may obscure the inference on the meson–nucleon interaction.

The differences between the square of the on-shell proton–proton scattering amplitude and the Jost function prescription are presented in figure 13a. To minimize the ambiguities that may result from

these discrepancies at least for the quantitative estimation of the effects of the unknown meson–nucleon interaction one can compare the spectra from the production of one meson to the spectra determined for the production of a meson whose interaction with nucleons is established. For instance, to estimate the strength of the ηpp and $\eta' pp$ FSI one can compare the appropriate observables to those of the $\pi^0 pp$ system.

Figures 18a and 18b show the dependence of $|M_0|$ on the phase-space volume for η and η' production normalized to $|M_0^{\pi^0}|$. The values of $|M_0|$ were extracted from the experimental data by means of equation (38) disregarding the proton–meson interaction ($|M_{p\eta(\eta') \rightarrow p\eta(\eta')}|$ was set to 1). If the influence of the neglected interactions were the same in case of $\eta(\eta')$ and π^0 the points would be consistent with the solid line. This is the case for the $pp \rightarrow pp\eta'$ reaction visualizing the weakness of the proton– η' interaction independently of the prescription used for the proton–proton FSI [37]. In case of the η' meson its low-energy interaction with the nucleons was expected to be very weak since there exists no baryonic resonance which would decay into $N\eta'$ [134]. Figure 18a shows — independently of the model used for the correction of the proton–proton FSI — the strong effects of the ηpp FSI at low V_{ps} .




Figure 18: The ratios of a) $|M_0^\eta|/|M_0^{\pi^0}|$ and b) $|M_0^{\eta'}|/|M_0^{\pi^0}|$ extracted from the data by means of equation (38), assuming a pp -FSI enhancement factor as depicted by the dotted line in figure 13a and neglecting the proton–meson interaction [37]. c) Arbitrarily normalized enhancement factors for pd -, $d\eta$ - and $p\eta$ -FSI. The η -proton factor is calculated according to equation (37), with $a_{p\eta} = 0.717 \text{ fm} + i 0.263 \text{ fm}$ [135] and $b_{p\eta} = -1.50 \text{ fm} - i 0.24 \text{ fm}$ [125]. The enhancement factor for η -deuteron has been extracted from the data of panel d) parametrizing the ratio of the cross section to the phase-space volume by the expression [136]: $F_{d\eta} = 1 + 0.5/(0.5 + (Q/5)^2)$. The proton–deuteron FSI factor is calculated according to ref. [70]. The figure is taken from [136]. d) Total cross section of the quasi-free $pn \rightarrow d\eta$ reaction as a function of the excess energy [48, 50]. The curve indicates the energy dependence proportional to the function $\sqrt{Q} \cdot (1 + Q/83.5)$. The relative s- and p-wave contribution was taken as determined for the $np \rightarrow d\pi^0$ reaction [43, 44, 47]. It was assumed that the p-wave to s-wave ratio is the same for the $pn \rightarrow d\eta$ and $pn \rightarrow d\pi^0$ reactions at a corresponding value of η_M .

With the up-to-date experimental accuracy, from all *Meson NN*-systems the ηNN one reveals by far the most interesting features. The dynamics of the ηNN system has become recently a subject of theoretical investigations in view of the possible existence of quasi-bound or resonant states [137]. According to reference [138] — within the present inaccuracy of $a_{\eta N}$ — the existence of quasi-bound η -mesic light nuclei could be possible. A direct measure of the formation — or non-formation — of an η -nuclear quasi-bound state is the real part of the η -nucleon scattering length [139]. The determined values of $\text{Re}(a_{\eta N})$ range between 0.20 fm [27] and 1.05 fm [144] depending on the analysis method and the reaction studied [122], and at present an univocal answer whether the attractive interaction between the η meson and nucleons is strong enough to form a quasi-bound state is not possible.

The shape of the energy dependence of the $pd \rightarrow {}^3\text{He} \eta$ cross section implies that either the real or imaginary part of the $\eta {}^3\text{He}$ scattering length has to be very large [140], which may be associated with a bound $\eta {}^3\text{He}$ system. Similarly encouraging are results of reference [141], where it is argued that a three-body ηNN resonant state, which may be formed close to the ηd threshold, may evolve into a quasi-bound state for $\text{Re}(a_{\eta N}) \geq 0.733$ fm. Also the close-to-threshold enhancement of the total cross section of the $pp \rightarrow pp\eta$ reaction was interpreted as being either a Borromean (quasi-bound) or a resonant ηpp state [142] provided that $\text{Re}(a_{\eta N}) \geq 0.7$ fm. Contrary, recent calculations performed within a three-body formalism [137] indicate that a formation of a three-body ηNN resonance state is not possible, independently of the ηN scattering parameters. Moreover, the authors of reference [143] exclude the possibility of the existence of an ηNN quasi-bound state. Results of both calculations [137, 143], although performed within a three-body formalism, have been based on the assumption of a separability of the two-body ηN and NN interactions. However, in the three-body system characterized by the pairwise attractive interactions, the particles can be pulled together so that their two-body potentials overlap, which may cause the appearance of qualitatively new features in the ηNN system [137]. Specifically interesting is the ηd final state where the pair of nucleons alone is bound by the strong interaction. Figure 18d shows the total cross section for the $pn \rightarrow d\eta$ reaction measured close to the production threshold. For excess energies below 10 MeV the data are enhanced over the energy dependence determined for the $pn \rightarrow d\pi^0$ reaction indicated by the solid curve. This is in qualitative agreement with the calculations of Ueda [145] for the three-body ηNN - πNN coupled system, which predict the existence of an ηNN quasi-bound state with a width of 20 MeV. Ueda pointed out that the binding of the ηNN system is due to the S_{11} ηN and 3S_1 $NN(d)$ interaction which is characterized by no centrifugal repulsion. Such repulsion makes the πNN system, in spite of the strong P_{33} πN attraction, hard to be bound [146]. Whether the observed cusp at the $pn \rightarrow d\eta$ threshold is large enough to confirm the existence of the ηNN bound state is recently vigorously discussed [147, 148]. The considerations concerning the possible existence of the light η -mesic nuclei will be continued in section 6.2.

The enhancement factor for the deuteron- η interaction inferred from the data in figure 18d varies much stronger in comparison to the proton- η one, as demonstrated in figure 18c. This suggests that the effects of this interaction should be even more pronounced in the differential distributions of the cross section for the $pd \rightarrow pd\eta$ reaction as those observed in case of the $pp \rightarrow pp\eta$ process, especially because the “screening” from the proton-deuteron interaction is by more than an order of magnitude smaller compared to the proton-proton interaction as can be deduced from the comparison of the solid lines in figures 18c and 13a. The experimental investigations on that issue [136, 149] as well as the search for the η -mesic nuclei [150] by measuring proton-deuteron induced reactions in the vicinity of the η production threshold are on the way.

4 Single meson production in NN scattering

4.1 Dynamics of pseudoscalar meson production

The considerations from the previous section led to the conclusion, that close to the kinematical threshold, the energy dependence of the total cross section is in the first approximation determined via the interaction among the outgoing particles and that the entire production dynamics manifests itself in a single constant, which determines the absolute scale of the total cross section. As a first step towards the understanding of the creation mechanism underlying the production let us compare the total cross sections of neutral pseudoscalar mesons π^0 , η and η' . Since the masses of these mesons are significantly different ¹² the influence of the kinematical flux factor F on the total cross section and the suppression due to the initial state interaction F_{ISI} depend substantially on the created meson. Therefore, for the comparison of the primary dynamics we will correct for these factors and instead of comparing the total cross section we will introduce — according to reference [37] — a dimensionless quantity $\sigma \cdot F/F_{ISI}$, which depends only on the primary production amplitude M_0 and on the final state interaction among the produced particles.




Figure 19: (a) Total cross section (σ) multiplied by the flux factor F and divided by the initial state interaction reduction factor F_{ISI} versus the available phase-space volume for the reactions $pp \rightarrow pp\eta$ (squares [50, 61–66]), $pp \rightarrow pp\pi^0$ (diamonds [8, 81, 89, 109, 151]), and $pp \rightarrow pp\eta'$ (circles [62, 90, 97, 111, 112]). (b) Experimentally determined ratio between $pn \rightarrow pn\eta$ and $pp \rightarrow pp\eta$ cross sections as a function of excess energy Q . The figure is adapted from reference [49].

Close-to-threshold the initial state interaction, which reduces the total cross section, is dominated by proton–proton scattering in the 3P_0 state which may be estimated in terms of phase-shifts and inelasticities by employing equation (42). The F_{ISI} factor is close to unity for pion production and amounts to ~ 0.2 [21] and ~ 0.33 [133] for the η and η' meson, respectively, at threshold.

A comparative study of the production of mesons with significantly different masses encounters the difficulty of finding a proper variable at which the observed yield can be compared. The total

¹²The π^0 , η and η' masses amount to 134.98 MeV/c², 547.30 MeV/c² and 957.78 MeV/c², respectively [134].

cross section is defined as the integral over the available phase-space volume of transition probabilities — reflecting the dynamics of the process — from the initial to the final state as written explicitly in equation 5. Thus, if the dynamics of the production process of two different mesons were exactly the same then the above introduced yield would also be strictly the same for both mesons provided it was extracted at the same value of V_{ps} . This inference would however not be valid if the production yields were compared at the η_M or Q variables. Therefore the volume of the available phase-space for the produced particles is the most suited quantity for the regarded comparison [37]. This could be also the best choice for the investigation of isospin breaking where the cross sections for the production of particles with different masses need to be compared (e.g. $\pi^+d \rightarrow pp\eta$ and $\pi^-d \rightarrow nn\eta$ [152]).

Figure 19a shows the yield of π^0 , η and η' mesons in the proton-proton interaction as a function of the available phase-space volume. The onset of higher partial waves is seen for π^0 and η mesons in the V_{ps} range between 10^3 and 10^4 MeV^2 , whereas the whole range covered by the η' data seems to be consistent with the pure Ss production. One can also recognize that the data for the Ss final state are grouped on parallel lines indicating a dependence according to the power law $\sigma \cdot F/F_{ISI} \approx \alpha \cdot V_{ps}^{0.61}$ [153] and that over the relevant range of V_{ps} the dynamics for η' meson production is about six times weaker than for the π^0 meson, which again is a further factor of six weaker than that of the η meson. This is an interesting observation, since the quark wave functions of η and η' comprise a similar amount of strangeness ($\approx 70\%$ [154]) and hence, in the nucleon-nucleon collision one would expect both these mesons to be produced much less copiously than the meson π^0 being predominantly built out of *up* and *down* quarks. On the hadronic level, however, one can qualitatively argue that the η meson owes its rich creation in the threshold nucleon-nucleon collisions to the existence of the baryonic resonance $N^*(1535)$ whose branching ratio into the $N\eta$ system amounts to 30–55% [134]. There is no such established resonance, which may decay into an s-wave $\eta'N$ system [134] and the π^0 meson production with the formation of the intermediate $\Delta(1232)$ state is strongly suppressed close-to-threshold, because of conservation laws.

Prior to the more comprehensive discussion of the production mechanism for different mesons let us present one more interesting observation visualizing that the production dynamics depends strongly not only on the structure of the created meson but also on the isospin configuration of the colliding nucleons.

For example, in the case of the η meson the comparison of the cross section from the production in proton-proton and proton-neutron collisions revealed that the yield of the η meson in the isosinglet nucleon configuration exceeds the one in the isospin triplet state by more than an order of magnitude. The ratio

$$R_\eta = \frac{\sigma(pn \rightarrow pn\eta)}{\sigma(pp \rightarrow pp\eta)},$$

presented in figure 19b, was determined to be about 6.5 in the excess energy range between 16 MeV and 109 MeV [49]. This implies that the production of the η meson with the total isospin equal to zero exceeds the production with isospin $I = 1$ by a factor of 12, since $\sigma(pn \rightarrow pn\eta) = (\sigma_{I=0} + \sigma_{I=1})/2$ and $\sigma(pp \rightarrow pp\eta) = \sigma_{I=1}$ and hence $\sigma_{I=0} = (2R_\eta - 1)\sigma_{I=1}$. On the mesonic level, this large difference of the total cross section between the isospin channels suggests the dominance of isovector meson (π and ρ) exchanges (figure 21 graph b) in the creation of η in nucleon-nucleon collisions [3, 49], but alternatively it can also be explained on the quark-gluon level assuming the instanton induced flavour-dependent quark-gluon interaction [24] (see below figure 26 graph e). The above examples visualize one of the most interesting problems in the investigations of the dynamics of the close-to-threshold meson production: Namely the determination of the relevant degrees of freedom for the description of the nucleon-nucleon interaction, especially in case when the nucleons are very close together¹³. As

¹³These investigations are listed as one of the key issues in hadronic physics [155].

pointed out by Nakayama [20], the transition region from the hadronic to constituent quark degrees of freedom does not have a well defined boundary and at present both approaches should be evaluated in order to test their relevance in the description of close-to-threshold meson production in the collision of nucleons.




Figure 20: A cartoon illustrating in naive geometrical terms that for $r < 2r_N + 2r_M$ meson exchange is unlikely to be appropriate to the description of the internucleon potential. Figure and caption are taken from reference [156].

Simple geometrical considerations presented in figure 20 indicate that at distances smaller than 2 fm the internucleon potential should begin to be free of meson exchange effects and may be dominated by the residual colour forces [156]. In the previous section it was shown, that the close-to-threshold production of mesons occurs when the colliding nucleons approach distances of about 0.5 fm in case of π^0 and of about 0.18 fm in case of ϕ production (see table 2). This distance is by about an order of magnitude smaller than 2 fm and it is rather difficult to imagine — in coordinate space — an exchange of mesons between nucleons as mechanism of the creation process. Such small collision parameters imply that the interacting nucleons — objects of about 1 fm — overlap and their internal degrees of freedom may be of importance.




Figure 21: Diagrams for the $pp \rightarrow pp \text{ Meson}$ reaction near threshold: (a) — *Meson-bremsstrahlung* (nucleonic current) (b) — “rescattering” term (nucleonic current) (c) — production via heavy-meson-exchange (d) — emission from virtual meson (mesonic current) (e) — excitation of an intermediate resonance (nucleon resonance current).

Nonetheless, in the last decade, the effective theory based on the meson exchanges, which accounts for the size of the participating particles by introduction of the momentum transfer dependent form factors, has been extensively utilized for the description of the creation process. Figure 21 represents the regarded mechanisms.

The question which processes — on the mesonic level — are responsible for the π^0 production was considered already in 1966 by Koltun and Reitan [6]. The authors deemed the direct pion production on one of the protons (fig. 21a) to play the most important role in the $pp \rightarrow pp\pi^0$ reaction. This supposition was anticipated until the precise experiments performed at IUCF [8,9] and CELSIUS [89] revealed that this mechanism can not account for more than 5 % of the total yield, as demonstrated in figure 22. Subsequent investigations showed that the inclusion of the processes illustrated in figure 21b, when the pion is produced on one of the protons and then scatters on the other one (in the πN s-wave [157] and πN p-wave via $\Delta(1232)$ [159]), increases the cross section essentially yet not enough to describe the data, even if the off-shell properties of the πN amplitude are taken into account [12,160]. The contribution of the rescattering mechanism is shown as the dashed line in figure 22. The agreement with the data can, however, be achieved — as shown by the solid line — if one takes additionally into account the mechanisms of heavy-meson-exchange associated with an intermediate virtual nucleon-antinucleon state [11,161] illustrated in figure 21c. A thorough discussion concerning π meson production in differently charged channels can be found e.g. in references [13,162].




Figure 22: Total cross section for the reaction $pp \rightarrow pp\pi^0$. The dotted line shows the results for direct production only. Adding rescattering gives the dashed line. Including also the contribution from heavy meson exchange leads to the solid line. The data are from refs. [8, 89]. The figure is adapted from reference [158]. Please note, that similarly as shown in figure 13b, the solid line representing the full calculation starts to deviate from the data at $\eta_M \approx 0.4$ ($Q \approx 11$ MeV), yet here the magnitude reflects the assumed mechanism and in figure 13 it was settled to fit the data. Thus indeed, the primary dynamics fixes in practice only the scale.

The dominance of the heavy-meson-exchange and the off-shell πN rescattering mechanisms, both reflecting the short range nucleon-nucleon interactions, confirms the simple estimation of the reaction region based on the momentum transfer between the colliding nucleons. Thus, in case of the π^0 meson creation, this approach succeeded to describe the magnitude of the total cross section. However, the meson-exchange model at the present stage does not reproduce the angular distributions of the spin observables for all $pp \rightarrow NN\pi$ isospin channels, simultaneously. Specifically, though the only available calculations of the spin correlation coefficients based on the Jülich model [14] are in a good agreement with the experimental data for the reactions $\vec{p}\vec{p} \rightarrow pn\pi^+$ [168] and $\vec{p}\vec{p} \rightarrow d\pi^+$ [169] they reveal serious discrepancies when confronted with the angular dependences of the spin correlation coefficients $A_\Sigma \equiv A_{xx} + A_{yy}$ and A_{zz} of the $\vec{p}\vec{p} \rightarrow pp\pi^0$ reaction [82]. Therefore, further investigations concerning the relative contributions of various mechanisms, as well as the different partial waves are still required.

The role of the mechanisms leading to the π production have been also considered in context of the lowest-order chiral perturbation theory [34] and relativistic impulse approximation [163]. In contrast to the meson exchange models the chiral perturbation method results [164–166] in the destructive interference between the direct and the pion-rescattering terms, and consequently implies more significant contributions originating from the shorter-than-pion-range mechanisms, like the heavy-meson exchanges or the creation of pions in the fusion of exchanged mesons [167].

In case of the η and η' mesons the investigations of the mechanisms underlying the production process are even more difficult. It is partly due to the fact that in contrary to the pion case even the coupling constants determining the strength of the $NN\eta(\eta')$ vertex are poorly established.

The values derived by various methods differ by more than an order of magnitude [154, 170]. For example, the estimation based on the dispersion relation results in $g_{NN\eta'}$ smaller than 1 [171], whereas the value obtained from the fits to low-energy nucleon-nucleon scattering in one-boson-exchange models amounts to 7.3 [172]. This uncertainty prevents from the exact estimation of the contribution from the nucleonic current. However, according to the systematic investigations of Nakayama et al. [20, 133, 173] this mechanism is of minor importance. The calculations have been performed with $g_{NN\eta'} = g_{NN\eta} = 6.1$ — used in the NN scattering analysis [174] — which is close to the upper limit of the predicted values.

It is at present rather well established [19, 33, 121, 173, 175–179] that the η meson is produced

predominantly via the excitation of the S_{11} baryonic resonance $N^*(1535)$ which subsequently decays into η and nucleon and whose creation is induced through the exchange of π , η , ρ , σ and ω mesons, as shown in figure 21e. Though all the quoted groups reproduce the magnitude of the total cross section, their models differ significantly as far as the relative contribution from the π , η and ρ exchange mechanisms is concerned. The discrepancies are due to the not well known strength of $Meson-N-S_{11}$ couplings and the ηN scattering potential. For example, while the dominance of the ρ meson exchange is anticipated by authors of references [33, 121, 175–177], it is rather the exchange of the η meson which dominates the production if one takes into account effects of the off-shell ηN scattering [180], or utilizes the multi-channel multi-resonance model [19]. In any case, the hitherto performed studies aiming to describe the total cross section show that the close-to-threshold production of η mesons in nucleon–nucleon collisions is dominated by the intermediate virtual S_{11} nucleon isobar whose width overlaps with the threshold. In order to disentangle between various scenarios of the S_{11} excitation a confrontation of the predictions with other observables is needed. The interference between considered amplitudes causes a different behaviour — depending on the assumed scenario — e.g. of the η meson angular distributions. These differences, however, are too weak in the close-to-threshold region to judge between different models. Also the ratio of the η meson production via the reactions $pp \rightarrow pp\eta$ and $pn \rightarrow pn\eta$ can be equally well described either by assuming the ρ meson exchange dominance [177] or by taking pseudoscalar and vector mesons for exciting the S_{11} resonance [173]. In the latter case, shown in figure 23, the excitation of the resonance via the ρ meson exchange was found to be negligible. Yet, the predictions of the analyzing power depends crucially on the assumed mechanism [173, 177]. This fact has triggered already experimental investigations which aim to determine the spin observables [181].




Figure 23: Total cross sections for the $pp \rightarrow pp\eta$ (upper panel) and $pn \rightarrow pn\eta$ (lower panel) reactions as a function of excess energy. The dashed curves correspond to the nucleonic current contribution and the dash-dotted curves to the mesonic current consisting of $\eta\rho\rho$, $\eta\omega\omega$ and $\eta a_0\pi$ contributions. The resonance current presented by the dotted line consists of the predominant $S_{11}(1535)$ and of the $P_{11}(1440)$ and $D_{13}(1520)$ resonances excited via exchange of π , η , ρ and ω mesons. The solid curves are the total contribution. The deviation from the data at low Q in the upper panel reflects the ηp FSI which was not included in the calculations. The data are from refs. [49, 61–65]. The figure is taken from [20].

Figure 23 presents also a contribution from the mesonic current where the η meson is created in the fusion of virtual e.g. ρ or ω mesons emitted from both colliding nucleons. In comparison to the

overwhelming strength of the resonance current this mechanism is however by a factor of 30 too weak. In contrast, it is enough to explain the magnitude of the close-to-threshold η' meson production in the proton–proton interaction, which is just by about a factor of 30 smaller compared to the η meson. Among the mesonic currents regarded by authors of reference [133] the $\rho\rho\eta'$ gives the dominant contribution, by a factor of five larger than the $\sigma\eta\eta'$ – and $\omega\omega\eta'$ –exchange. However, the understanding of the η' production on the hadronic level is far from being satisfactory. The magnitude of the total cross section was also well reproduced in the frame of a one–boson–exchange model (nucleonic current) where the virtual boson ($B = \pi, \eta, \sigma, \rho, \omega, a_0$) created on one of the colliding protons converts to the η' on the other one [35]. Taking into account the off–shell effects of the $Bp \rightarrow \eta'p$ amplitude it was found that the short range σ and ρ meson exchanges dominate the creation process, whereas the π exchange plays a minor role. On the contrary, other authors [182] reproduced the magnitude of the total cross section with π exchange only and concluded that the η, ρ and ω exchange currents either play no role or cancel each other. However, in both of the above quoted calculations [35, 182] the initial state interaction between protons, which reduces the rate by a factor of about 3, was not taken into account and hence the obtained results would in any case reproduce only 30 % of the entire magnitude and could be at least qualitatively reconciled with the mentioned result of reference [133], where the nucleonic current was found to be small. Moreover, the choice of the another prescription for the form factors could reduce the one–pion–exchange contribution substantially [133]. However, the picture that the η' meson is predominantly created through the mesonic current, which we would like to advocate, remains at present unclear as well. This is because the magnitude of the total cross section could have been also described assuming that the production of η' is resonant [133]. As possible intermediary resonances the recently reported [183] $S_{11}(1897)$ and $P_{11}(1986)$ have been considered. These resonances were deduced from η' photoproduction data, under the assumption that the close-to-threshold enhancement observed for the $\gamma p \rightarrow \eta'p$ reaction can be utterly assigned to resonance production. Further, as well strong, assumptions have been made in the derivation of the $g_{NN^*\eta'}$ and $g_{NN^*\pi}$ coupling constants [133]. Hence, it is rather fair to state that in the case of the close-to-threshold π^0 and η meson production in nucleon–nucleon collisions the dynamics is roughly understood on the hadronic level, but the mechanisms leading to the η' creation are still relatively unknown. Until now it was not possible to estimate satisfactorily the relative contributions of the nucleonic, mesonic and resonance current to the production process. In fact, model uncertainties allow that each one separately could describe the absolute values of the $pp \rightarrow pp\eta'$ total cross section. This rather pessimistic conclusion calls for further experimental and theoretical research. The understanding of the production dynamics of the η' meson on both hadronic and the quark–gluon level is particularly important since its wave function comprises a significant gluonic component [184], distinguishing it from other mesons and hence the comprehension of the mechanism leading to its creation in collisions of hadrons may help in the determination of its quark–gluon structure which is also of importance to investigate possible glueball candidates [184]. Hereafter we will briefly report about the gluonic mechanisms which may — additionally to the meson exchange processes discussed above — contribute to the η' production.

As already mentioned, the close-to-threshold production of η and η' mesons in the nucleon–nucleon interaction requires a large momentum transfer between the nucleons and hence can occur only at distances of about 0.3 fm (see table 2). This suggests that the quark–gluon degrees of freedom may indeed play a significant role in the production dynamics of these mesons. A possibly large glue content of the η' and the dominant flavour–singlet combination of its quark wave function may cause that the dynamics of its production process in nucleon–nucleon collisions is significantly different from that responsible for the production of other mesons. In particular, the η' meson can be efficiently created via a “contact interaction” from the glue which is excited in the interaction region of the colliding nucleons [185]. A gluon–induced contact interaction contributing to the close-to-threshold $pp \rightarrow pp\eta'$ reaction derived in the frame of the U(1)–anomaly extended chiral Lagrangian is discussed in references [185–187]. The

strength of this contact term is related to the amount of spin carried by polarized gluons in a polarized proton [187,188], thus making the study of the close-to-threshold η' meson production even more interesting.

Figure 24 depicts possible short-range mechanisms which may lead to the creation of the η' meson via a fusion of gluons emitted from the exchanged quarks of the colliding protons [189] or via an exchange of a colour-singlet object made up from glue, which then re-scatters and converts into η' [191]. The hadronization of gluons to the η' meson may proceed directly via its gluonic component or through its overwhelming flavour-singlet admixture η_1 (see fig. 25). Contrary to the significant meson exchange mechanisms and the fusion of gluons of figure 24 graph a), the creation through the colour-singlet object proposed by S.D. Bass (graph 24b) is isospin independent, and hence should lead to the same production yield of the η' meson in both reactions ($pp \rightarrow pp\eta'$ and $pn \rightarrow pn\eta'$) because gluons do not distinguish between flavours. This property should allow to test the relevance of a short range gluonic term [192] by the experimental determination of the cross section ratio $R_{\eta'} = \sigma(pn \rightarrow pn\eta')/\sigma(pp \rightarrow pp\eta')$, which in that case should be close to unity after correcting for the final and initial state interaction between participating baryons. The other extreme scenario — assuming the dominance of the isovector meson exchange mechanism — should result in the value of $R_{\eta'}$ close to 6.5 as was already established in the case of the η meson (see fig. 19b). The experimental investigations of that issue are in preparation [51].




Figure 24: Diagrams depicting possible quark-gluon dynamics of the reaction $pp \rightarrow pp\eta'$. (a) — production via a fusion of gluons [189] with rearrangement of quarks. (b) — production via a rescattering of a “low energy pomeron”.




Figure 25: Coupling of η and η' to two gluons through (a) quark and antiquark triangle loop and (b) gluonic admixture. η_1 denotes the flavour-singlet quark-antiquark state. This indicates that gluons may convert into the η or η' meson via a triangle quark loop only by coupling through their flavour singlet part. The figure is taken from reference [190].

In addition to the QCD motivated investigation presented above [185,186], the interesting features of the close-to-threshold meson production, in particular the large cross section for the η meson in proton-proton interactions exceeding the one of the pion, or even more surprisingly large cross section of the η production in proton-neutron collisions, encouraged also other authors to seek for the underlying — OZI rule violating — creation mechanisms in the frame of microscopic models of QCD [24, 25, 193]. The hitherto regarded processes are presented in figure 26. As indicated in the pictures the structure of participating baryons has been modeled as quark-diquark objects with harmonic confinement [194]. In the upper graphs the large momentum transfer is shared by the exchanged gluons with a subsequent interchange of quarks to provide a colourless object in the final state. The lower graphs depict the two examples of instanton induced interactions with a 6-quark-antiquark (fig. 26d) and two-gluon vertex (fig. 26e).

Adjusting the normalization to the cross section of the $pp \rightarrow pp\pi^0$ reaction at a single energy point the model [25] accounts roughly for close-to-threshold cross sections of other pseudoscalar and vector mesons in proton-proton collisions. Though this approach is characterized by significantly smaller number of parameters than in the meson exchange models, their uncertainties allow for the description of the data equally well either by the gluon exchange or by the instanton induced interactions, at least in case of the π^0 , η' , ω and ϕ mesons. Yet for the resonance dominated η and K^+ meson production in proton-proton collisions it was found that the instanton induced interaction presented by graph 26d is not sufficient. Similarly, the authors of reference [24] argue that the instanton induced interaction with a quark-gluon vertex (graph 26e) should be of no importance for the η production in proton-proton collisions. This arises from the properties of the vertex which lead to the η production only in case of an interaction between quarks of different flavours ($ud \rightarrow udg$) and in the quark-diquark model of baryons the proton consists of ud -diquark and u -quark and correspondingly the neutron is modeled as ud -diquark and d -quark. Although negligible in case of proton-proton interactions, this mechanism may contribute significantly to the total cross section of the $pn \rightarrow pn\eta$ reaction and indeed as shown in reference [24] it reproduces the data.




Figure 26: Diagrams for the $qq \rightarrow qq(q\bar{q})$ production operator: (a), (b) Two-gluon exchange and rescattering mechanism (c) Correlated, colourless two-gluon exchange. The dashed line indicates the excitation of an intermediate NN^* system. (d) Instanton induced 6-quark interaction. Figures (a–d) according to [25]. (e) The instanton contribution to the η meson production in the proton–neutron interaction. Figure copied from [24].

However, the magnitude of the cross section is very sensitive to the size of the instanton in the QCD vacuum and — similar to the uncertainty of coupling constants in case of the hadronic approach — at present it makes precise predictions impossible. Thus, despite the partial successes, at the present stage of developments both approaches — on the quark–gluon and on the hadronic level — do not deliver an unambiguous answer to the dynamics of the close-to-threshold meson production in the nucleon–nucleon interaction.

4.2 Vector meson production

Vector meson production studies in the mass range up to $1 \text{ GeV}/c^2$ in hadronic collisions are especially motivated by questions concerning the strangeness content of the nucleon. Within the quark model the ϕ meson is an (almost) ideally mixed $s\bar{s}$ state, whereas the nucleon is composed only of u - and d -quarks. According to the OZI rule [195]¹⁴, ϕ production in the nucleon–nucleon interaction would be forbidden as a process with disconnected quark lines, if the ϕ was a pure $s\bar{s}$ state.

However, due to a small deviation from ideal mixing ($\delta_V = 3.7^\circ$ [134]) the ϕ meson is allowed to couple to the nucleon through its $(u\bar{u} + d\bar{d})$ admixture. Quantitatively, the naive OZI rule results in a

¹⁴for a review see [196].

suppression of ϕ compared to ω production in hadronic interactions

$$R = \frac{\sigma(A B \rightarrow \phi X)}{\sigma(A B \rightarrow \omega X)} = \tan^2(\delta_V) \approx 4.2 \times 10^{-3} \quad (43)$$

after phase-space corrections, where A , B and X denote hadronic systems consisting only of light quarks [197]. Any significant deviation might be interpreted as a hint for a strangeness component in the nucleon. A considerable contribution of strange sea quarks to the nucleon wave function is suggested by results on the $\Sigma_{\pi N}$ term in pion nucleon elastic scattering [198], on the nucleon's structure function in deep inelastic scattering with polarized muons [199] and, more recently, on charm production in deep inelastic neutrino scattering [200, 201].

Large violations of the OZI expectation (eq. (43)) have been found in $p\bar{p}$ annihilation experiments by the ASTERIX, Crystal Barrel and OBELIX collaborations at LEAR (as reviewed in [202]). However, deviations are predominantly found in $p\bar{p}$ annihilation at rest as compared to higher energies and seem to be restricted to S-wave processes only. Furthermore, the results are strongly dependent on the final state: While the $\phi\pi$ and $\phi\gamma$ channels exceed the OZI value by a factor of 20 and 100, respectively, little or no effect is seen in the $\phi\pi\pi$ and $\phi\eta$ final states.

These data have been interpreted as “shake-out” and “re-arrangement” of a negatively polarized $s\bar{s}$ Fock space component of the proton wave function [202, 203], which also accounts for large double $\phi\phi$ cross sections [204] in $p\bar{p}$ annihilation measured at the JETSET experiment [205]. However, OZI-allowed two-step processes via intermediate $K\bar{K}$ or $\Lambda\bar{\Lambda}$ states have been shown to describe available data from $p\bar{p}$ annihilation without any strangeness in the nucleon [206–213].

In nucleon–nucleon scattering, effects of competing two step processes for ϕ production via intermediate ΛK or ΣK states are expected to be of minor importance [212, 214]. Furthermore, in close-to-threshold ϕ production the entrance channel must be in a 3P_1 state (see table 1) due to parity and angular momentum conservation and it is the spin triplet fraction in $p\bar{p}$ annihilation which is strongly correlated to the ϕ meson yield [215]. Thus, cross section ratios for ϕ and ω production in proton–proton scattering should clearly indicate possible OZI violations and probe the $s\bar{s}$ component of the nucleon.

The exclusive production ratio has been determined in the reaction $pp \rightarrow pp\phi/\omega$ at a beam momentum of 3.67 GeV/c corresponding to excess energies of 82 MeV and 319 MeV to the ϕ and ω thresholds, respectively, by the DISTO collaboration at SATURNE [216]. After phase-space corrections, the observed ratio exceeds the OZI expectation by an order of magnitude, while data at excess energies larger than 1.1 GeV [217] show an enhancement smaller by a factor of four (see fig. 27). One-boson-exchange models [218–221] underestimate the absolute value of the ϕ production cross section measured at the DISTO experiment by at least a factor of three.




Figure 27: ϕ/ω total cross section ratio in proton–proton collisions versus excess energy with respect to the ϕ production threshold from [216]. The recent data point from the DISTO collaboration (solid square [216]) is shown together with high energy data (solid circles [217]) and calculations taking into account the OZI prediction and three–body phase–space [216] and results obtained within a one–boson–exchange model including the proton–proton FSI [218]. Data includes both statistical and systematical errors.

Figure 27 includes results obtained within a one–pion–exchange model [218], which provides a reasonable description of high energy data. Model predictions — taking into account the proton–proton final state interaction only — falls short of the recent close–to–threshold value [216]. Using instead a parametrization of the energy dependence of the ω production cross section and considering both the finite ω width as well as effects of proton–proton FSI [219, 222], the ϕ/ω ratio can be determined at equal excess energies, thus reducing uncertainties due to the available phase–space, partial wave amplitudes ¹⁵ and effects of the proton–proton FSI. In conclusion, the excess over the OZI prediction is decreased to a factor of five [216], i.e. to a modest enhancement as compared to the results from $p\bar{p}$ annihilation experiments ¹⁶.

In the DISTO results, ϕ production at an excess energy of 82 MeV appears to be dominated by s–wave relative to the nucleon–nucleon system in agreement with theoretical predictions [224] and phenomenologically described in section 3. However, the proton–proton angular distribution relative to the ϕ direction is found to deviate significantly from isotropic emission at this beam energy with evidence for a P–wave contribution in the proton–proton subsystem. Consequently, ϕ production at 82 MeV above threshold, although dominated by a 3P_1 proton–proton entrance channel, partly proceeds via the 1S_0 and 1D_2 spin singlet partial waves [216]. Thus, the influence of the spin–triplet entrance channel, prerequisite to a “re–arrangement” of an $s\bar{s}$ component in the nucleon wave function [202], might be diluted in the presently available data and would be expected to be more prominent in data closer to the production threshold.

¹⁵The ω angular distribution relative to the pp system measured at an excess energy of 319 MeV shows evidence for p– and d–waves contributing [216].

¹⁶Comparing the DISTO data at $Q = 82$ MeV [216] and the recent COSY–TOF result at an excess energy of 92 MeV [223] leads to a similar enhancement of ≈ 7 compared to the OZI expectation.




Figure 28: Vector meson production currents in nucleon–nucleon scattering ($NN \rightarrow NNv$) studied within a relativistic meson exchange model in [20, 225, 226]:

(a) nucleonic current and (b) mesonic current.

$M = \pi, \eta, \rho, \omega, \sigma, a_0$ denotes the exchanged meson and B the intermediate off mass-shell baryon.

Recently, within the framework of a relativistic meson exchange model, nucleonic and meson–exchange currents have been used explicitly to study vector meson production in nucleon–nucleon scattering (fig. 28 [20, 225, 226]). Uncertainties in the model calculation for the nucleonic current arise mainly from a large range of values quoted for vector and tensor NNv coupling constants [171, 174, 212, 227–232], as well as from the vector and tensor vertex form factors for the coupling of an on–mass shell vector meson to off mass–shell nucleons. Note, that a resonance contribution to the nucleonic current (fig. 28a) is neglected in the model [225], since there are no isospin–1/2 nucleon resonances experimentally known coupling to the vp channel [233]. Due to the strength of both the $NN\pi$ and $NN\rho$ couplings and the $g_{\pi\rho\omega}$ coupling constant as compared to other possible choices, only the $\pi\rho v$ vertex is considered in evaluating the meson–exchange current. For ω production, contributions from the latter and the nucleonic current are found to interfere destructively when added coherently, which leaves two choices for a given value of the $NN\omega$ tensor–to–vector coupling ratio — when fitting the close–to–threshold data on the reaction $pp \rightarrow pp\omega$ taken at SATURNE [236] — for the only remaining free parameter of the model, the cut–off parameter at the $NN\omega$ form factor. Effects from a destructive interference of mesonic exchange current and nucleonic current contribution are also concluded by other authors for ω production in [234]¹⁷ and for ϕ production in [235]. In Nakayama’s analysis the energy dependence of the total cross section is slightly influenced by either choice, as shown by the solid and dashed curves¹⁸ in figure 29 together with a parametrization based on one–pion–exchange (dash–dotted line [218]). In comparison, total cross section data taken at the SPES 3 spectrometer up to excess energies of 30 MeV [236], the DISTO data point discussed above [216], high energy data from [238] and recent results obtained at the TOF facility at COSY at excess energies of $Q = 92$ MeV and 173 MeV, respectively [223], are included.

¹⁷Kaiser’s study [234] is based on the calculation of various tree level diagrams in comparison with the threshold transition amplitude extracted from data of [236] and follows the approach developed in [34, 237] (see section 4.4). The dominant nucleonic current contribution arises from ρ^0 exchange. However, the combined result of the studied amplitudes shows similarly to [225, 226] a dependence on the choice of the tensor–to–vector coupling ratio.

¹⁸The two curves correspond to the extreme cases found for the tensor coupling values $f_{NNv} = \pm 0.5 g_{NNv}$ [226], with cut–off masses of 1545 MeV (dashed line) and 1170 MeV (solid line), respectively.




Figure 29: Total cross section for $pp \rightarrow pp\omega$ as a function of the excess energy. Data are from refs. [216] (open triangle), [223] (solid squares), [236] (solid circles) and [238] (open squares). Calculations include a one-pion-exchange based parametrization (dash-dotted curve [218]) and results obtained within a relativistic meson-exchange model considering nucleonic and mesonic current contributions (solid and dashed curves [225, 226], for details see text). The figure is taken from ref. [223].

The one-pion-exchange parametrization [218] deviates strongly from the close-to-threshold data, where FSI effects become important, which have not been included in the calculation. Both predictions obtained within the relativistic meson exchange model of [20, 225, 226] overestimate the recent COSY-TOF data [223] by at least a factor of five. The solid curve in figure 29 seems to follow the data slightly better, which would correspond to a lower cut-off parameter in the nucleonic current and, consequently, a dominating meson exchange current contribution.

Within the model [225, 226], angular distributions of the produced mesons are expected to differ depending on the dominant production mechanism for both ω and ϕ production: While the mesonic current leads to an isotropic angular distribution, the nucleonic current results in a $\cos^2 \Theta_v$ behaviour for the vector meson emission angle Θ_v in the overall centre-of-mass system¹⁹. Thus, angular distributions offer a unique possibility to separate both contributions, or to isolate the nucleonic current contribution in order to determine the ϕNN coupling directly in a combined analysis of ω and ϕ production data in pp induced reactions: The experimentally observed isotropic ϕ angular distribution [216] clearly favours a dominant $\phi\rho\pi$ meson exchange over the nucleonic current [20, 212]. Contrary, the centre-of-mass angular distribution of the ω meson obtained at the COSY-TOF facility at an excess energy of 173 MeV [223] exhibits a strong anisotropy similar to the observation by the DISTO collaboration at an excess energy of 319 MeV [216], indicating the presence of higher partial waves and rather favours a dominant nucleonic current contribution instead of meson exchange currents suggested by the shape of the energy dependence of the total cross section in figure 29²⁰.

Further constraints on the relevant vector meson production mechanisms in nucleon-nucleon scattering might be obtained from comparing with complementary meson induced reactions: An analysis of $\pi N \rightarrow N\phi/\omega$ data concludes ω production to result from strong interferences between the meson exchange current and both nucleonic and resonance currents, which contribute with similar strengths [240]. Near threshold, the P_{11} $N^*(1440)$ resonance and the S_{11} $N^*(1535)$ and $N^*(1650)$ excitations are found to contribute, with the S_{11} contributions canceling to a large extent²¹.

In conclusion, at least for the reaction $\pi N \rightarrow N\omega/\phi$, the nucleonic current contribution appears to

¹⁹The $\cos^2 \Theta_v$ behaviour arises according to [225] from the spin-dependent part of the nucleonic current contribution, the “magnetization current” [239].

²⁰It should be noted, that the excess energy of the COSY-TOF data [223] is slightly out of the energy range covered by the predictions from Nakayama et al. [225, 226] due to the onset of inelasticities in the final state.

²¹For the resonance contributions in the work of Titov et al. [240], ωNN^* couplings and relative phases have been taken from [241].

be much larger for ω production relative to the meson exchange current, contrary to ϕ production [240]. Thus, the scenario from pion induced vector meson production seems to agree — qualitatively — with the observations in nucleon–nucleon scattering, and is interesting to note especially in view of the ω angular distribution from the COSY–TOF data [223]. However, angular distributions at lower excess energies would clearly help to constrain the interpretation of the presently available nucleon–nucleon data in future.

In a combined analysis of ω and ϕ production within the relativistic meson exchange framework [20] a value of the ϕ coupling to the nucleon has been extracted from the DISTO data and is found to be consistent with the OZI expectation $g_{NN\phi} = -3 g_{NN\rho} \sin \delta_V \approx -(0.60 \pm 0.15)$ [226]. Yet, as stated by Nakayama, the value still has large uncertainties, especially due to the lack of more and precise data in the threshold region [20]. Indeed, on the basis of extracted transition matrix elements [219] combining πN and pp induced reactions, a ϕ/ω production ratio has been derived, which exceeds the OZI estimate of eq. (43) by a factor of five in both cases.

As suggested in [216], future experiments on ϕ and ω production in close-to-threshold proton–neutron collisions might provide further evidence for the intrinsic strangeness content of the nucleon, as predictions for ϕ production cross sections based on the “shake-out” and “rearrangement” mechanism [242] and on meson exchange models [221, 224] differ by more than an order of magnitude.

4.3 Scalar sector

The 1 GeV/c² meson mass range is continuously under discussion regarding the nature of the scalar resonances $f_0(980)$ and $a_0(980)$, which have been interpreted as exotic two–quark two–antiquark states [243], possible glueball candidates [244], excitations of the light quark vacuum condensate (*minions* [245]), conventional $q\bar{q}$ states [246, 247], or molecular like $K\bar{K}$ bound states [248, 249]. Just already the similarity of the masses of these objects with twice the kaon mass, see figure 30a, suggests a strong correlation between these resonances and the $K\bar{K}$ system.

Within the framework of the Jülich meson exchange model for $\pi\pi$ [251] and $\pi\eta$ [252] scattering the $K\bar{K}$ interaction dominated by vector meson exchange gives rise to a $K\bar{K}$ bound state identified with the $f_0(980)$ in the isoscalar sector, while the isovector $a_0(980)$ is concluded to be a dynamically generated threshold effect [252, 253].

As shown in figure 30b, both shape and absolute scale of $\pi\pi \rightarrow K\bar{K}$ transitions turn out to depend crucially on the strength of the $K\bar{K}$ interaction, which, in turn is prerequisite of a $K\bar{K}$ bound state interpretation of the $f_0(980)$. Similar effects might be expected for the elementary kaon–antikaon production in proton–proton scattering.

Although the $K\bar{K}$ decay mode of $f_0(980)$ and $a_0(980)$ is rather weak in comparison to the dominant $\pi\pi$ and $\pi\eta$ decay channels [134], even a new theoretical analysis based on the chiral approach can not account for the $f_0(980)$ and $a_0(980)$ if the $K\bar{K}$ channel is not introduced additionally to the $\pi\pi$ and $\pi\eta$ interaction [254]. An analysis of the $\pi\pi$ and $K\bar{K}$ interaction [255] showed that the f_0 corresponds to two poles on the second and third Riemann sheet, respectively, and appears physically as an object with a decay width of about 400 MeV and a narrow peak width of about 50 MeV. The same parameters of the f_0 were found by utilizing a unitarized quark model, according to which the f_0 was interpreted as a $q\bar{q}$ state with a large admixture of a $K\bar{K}$ virtual state [256]. The origin of the scalar resonances was also thoroughly studied by means of a coupled channel analysis considering $\pi\pi$, $K\bar{K}$ and $\sigma\sigma$ meson–meson scattering [257, 258]. Decreasing gradually the interchannel coupling constants it was inferred that for some solutions at the limit of the fully uncoupled case the f_0 corresponds to a $K\bar{K}$ bound state [258, 259].

Recently notable progress has been made concerning the discussion of this topic due to approaches in unitary extensions of chiral perturbation theory where the resonances under consideration qualify as dynamically generated resonances from the multiple scattering with the lowest order chiral Lagrangian,

as outlined in the review [260].




Figure 30: (a) Weighted mass average for the $f_0(980)$ scalar resonance from [250]. (b) Predictions of the Jülich model for $\pi\pi \rightarrow K\bar{K}$ transitions using the full model and neglecting any $K\bar{K}$ interaction, taken from ref. [252].

In high energy experiments, the f_0 meson is observed as a resonance in the system of two pions produced in a variety of hadro-production reactions [261–264] and in the hadronic decays of either heavier mesons [265–267] or the Z^0 boson [268,269] created in e^+e^- collisions. These experiments study the invariant masses of the created neutral ($\pi^0\pi^0$) [261,263,267] and charged ($\pi^+\pi^-$) [262,264–266,268] pion pairs. Similarly, charged [270,271] and neutral a_0 [272] mesons were observed as a clear signal in an invariant mass spectrum of the $\eta\pi$ system.

Complementary to these approaches, studying the interaction of $\pi\pi$, $K\bar{K}$ and $\pi\eta$ meson pairs, the COSY-11 collaboration presently investigates the possible manifestation of the mesons f_0 and/or a_0 as doorway states leading to meson production in proton-proton collisions, namely $pp \rightarrow pp f_0(a_0) \rightarrow pp \text{ Mesons}$. By measuring the missing mass of the pp -system the overlapping f_0 - a_0 resonances are studied as a genuine particle produced directly at the reaction. The production of these resonances is carried out at a mass range a few tens of MeV below the $K\bar{K}$ threshold (where the resonances can only decay into non-strange mesons) and above the $K\bar{K}$ thresholds, where the equivalent part of these broad resonances can be excited.

It is obvious that for the production of a broad resonance the phrase “close-to-threshold” is not well defined and implies here that the beam momentum is such that masses just in the range of the resonance can be excited. Different mass ranges of such objects are produced with different excess energies making the interpretation of the experimental data difficult but still possible [278]. It is worth noting, that recently first measurements of the f_0 meson production relatively close to its threshold but still above the $K\bar{K}$ threshold were performed by observing the ϕ meson decays into the $f_0\gamma$ channel [266,267], where the ϕ is only about 40 MeV/ c^2 heavier than the f_0 . In a recent publication [77] data on the close-to-threshold K^+K^- production following the proton-proton interaction at an excess energy of $Q = 17$ MeV are presented. The obtained distribution of the missing mass to the pp system is shown in figure 31 and demonstrates that the non-resonant K^+K^- production (solid line) is hardly distinguishable from the resonant $pp \rightarrow pp f_0(980) \rightarrow pp K^+K^-$ reaction sequence (dashed line) [273].




Figure 31: Experimental spectrum of the K^+K^- invariant mass measured for the reaction $pp \rightarrow ppK^+K^-$ at a beam momentum of 3.356 GeV/c corresponding to an excess energy of $Q = 17$ MeV with respect to the K^+K^- threshold (data points). The width of the bins corresponds to the experimental resolution of the mass determination (FWHM ≈ 2 MeV). The solid and dashed lines show Monte-Carlo simulations assuming the direct and resonant production, correspondingly [273].

Since the statistics of the data were not sufficient to favour one of the two processes, the cross section was extracted for both and resulted in nearly identical values of: $\sigma_{non-resonant} = (1.80 \pm 0.27^{+0.28}_{-0.35})$ nb and $\sigma_{resonant} = (1.84 \pm 0.29^{+0.25}_{-0.33})$ nb including statistical and systematical errors, respectively. The issue, whether there is a chance to distinguish between $K\bar{K}$ pairs originating from the decay of a genuine f_0/a_0 resonance and those produced by a strong $\pi\pi-K\bar{K}$ correlation is at present under theoretical investigation [253]. Still, calculations based upon the one-pion exchange and a Breit-Wigner presentation of the f_0 resonance indicate [274] that no f_0 signal might be extracted from the K^+K^- invariant mass spectrum in $pp \rightarrow ppK^+K^-$ reaction at near-threshold energies, due to the large contribution from other reaction channels [79] as arising from pion and kaon exchange.

Considering the $a_0(980)$ as a conventional $q\bar{q}$ meson, predictions on the total and differential cross sections for a_0 production via the reaction $pp \rightarrow da_0^+$ are given for the COSY energy range in [275]: Within a two-step model, where the a_0 is found to be produced predominantly from u-channel π exchange with a subsequent fusion of the nucleons to a deuteron bound state, a rather large total cross section is concluded. An experimental verification at the ANKE facility [276] might give an additional hint on the $q\bar{q}$ content of the isovector scalar resonance. The measurements also aim at determining the rather poorly known a_0 branching ratios to the $K\bar{K}$ and $\eta\pi$ final states (see discussion in [275] and references therein).

The role of final state interactions in the reactions $pp \rightarrow dK^+\bar{K}^0$ and $pp \rightarrow d\pi^+\eta$ is explicitly studied in [277]: Considering primary production amplitudes within meson-baryon chiral perturbation theory, $\bar{K}^0d$ final state interactions — due to the proximity of the $\Lambda(1405)$ to the $\bar{K}^0n$ system — are expected to change the cross section by up to an order of magnitude and to significantly influence invariant mass distributions. The $K^+\bar{K}^0$ and $\pi^+\eta$ final state systems are investigated within a coupled-channel chiral unitary approach, in which the $a_0(980)$ is generated dynamically as a meson-meson resonance. Once the two remaining independent model parameters — related to the couplings of the two pseudoscalar meson systems — are fixed by sensitive experimental data, i.e. $K^+\bar{K}^0$ and $\bar{K}^0d$ invariant mass distributions, the approach will provide absolute predictions for the $\eta\pi$ system. Thus, the underlying dynamics of the coupled-channel system giving rise to the $a_0(980)$ should be probed directly by the experimental approach of [276].

Further experimental efforts on studying the iso-singlet f_0 and the iso-triplet a_0 scalar resonances are continuing at the COSY-11 [278,279] and ANKE [280] facilities at COSY using different approaches.

4.4 Associated strangeness production

In elementary hadronic interactions with no strange valence quark in the initial state the associated strangeness production provides a powerful tool to study reaction dynamics by introducing an “impurity” to hadronic matter. Thus, quark model descriptions might be related to mesonic or baryonic degrees of freedom, with the onset of quark degrees of freedom expected for kinematical situations with large enough transverse momentum transfer.

A large fraction of the nucleon spin is carried by the intrinsic orbital angular momentum of quarks which – in the condensate – are pairwise in a 3P_0 state. The challenge of the strange quark, with its intermediate mass between light and heavy quarks, is to explore the effective degrees of freedom via the associated strangeness production in the low energy structure of QCD. The $\bar{q}q$ ($\bar{s}s$) vacuum condensate is the link between current quark and gluon of QCD and valence quark and potentials of the quark model, structures which can give the massless QCD quarks their constituent masses.

In close-to-threshold kaon-hyperon production, effects of the low energy hyperon-nucleon interaction are inherent to the observables, and allow to constrain hyperon nucleon interaction models.

As the strange quark is heavy compared to up and down quarks ($m_u, m_d \ll m_s \sim \Lambda_{QCD}$) it is not a priori clear, whether the strange quark is appropriately treated as a light quark on the hadronic scale in chiral perturbative approaches with an expansion parameter of $m_K/4\pi F_\pi = 0.43$ (with the kaon mass m_K and the pion decay constant F_π) compared to $m_\pi/4\pi F_\pi = 0.12$. In exploratory calculations, SU(3) baryon chiral perturbation theory only has been applied to kaon photo- and electroproduction [281], for a more detailed discussion see [282]).

With a significant contribution of strange sea quarks to the nucleon wave function being suggested by complementary experimental approaches during the past years (see section 4.2 and [198–201]) the creation of an $s\bar{s}$ pair in the nucleon-nucleon interaction should provide a further piece of valuable information on the structure of the nucleon [193]. Especially polarization observables are expected to act as very sensitive tools [283].

At the same time, both the kaon and antikaon (see section 5.4) production cross section in nucleon-nucleon scattering are important input parameters within the framework of transport model calculations used to describe the strangeness flow in heavy ion collisions: Due to strangeness conservation, kaons will not be absorbed in the nuclear medium. Thus, kaon yields in subthreshold production are expected to probe the early stage of the collision, i.e. hot, dense matter, the nuclear equation of state at high densities [284] and the in-medium kaon potential [285–287], see also section 5.4), which relates to the partial restoration of chiral symmetry [288,289] (for a review see [290]). Elementary cross sections are also of interest in view of the recent inclusive subthreshold measurements in nucleus-nucleus collisions, resulting in comparable K^+ and K^- yields at the same energy per nucleon below threshold [291–293].

Exclusive data close-to-threshold on the elementary kaon production via $pp \rightarrow pK^+\Lambda$ have only become available recently with the advent of dedicated experiments at COSY. Figure 32 shows the available total cross section data obtained at the COSY-11 [113–115] and COSY-TOF facility [116–118] together with an earlier result from the BNL Cosmotron [119] and theoretical calculations and parametrizations as a function of the excess energy Q .




Figure 32: Total cross section of the reaction $pp \rightarrow pK^+\Lambda$. Data obtained at the COSY-11 facility are marked by filled [113,114] and open squares [115], COSY-TOF data by filled circles [116] and triangles [117,118] and the BNL result [119] by an open cross, respectively. Experimental data are the same as in figure 14b. The arrow denotes the opening of the $pK^+\Sigma^0$ threshold. For comparison, theoretical descriptions [294–300] are included (for details see text).

Assuming the production amplitude to be constant and both the $p\Lambda$ system and the K^+ relative to the $p\Lambda$ subsystem to be in s-wave leads to a production cross section proportional to η^4 (section 2.1, eq. (16)). This dependence has been adopted by Schürmann and Zwermann [294] for a quartic (in terms of the maximum meson momentum) parametrization of the cross section (black solid line in fig. 32), leading to a reasonable fit [295] of data at excess energies above 150 MeV. Although quite commonly employed in the description of nucleon–nucleus and nucleus–nucleus collisions in the past [297], the quartic parametrization underestimates the close-to-threshold data by an order of magnitude, as does a linear parametrization once suggested by Randrup and Ko (grey solid line [296]), overestimating the presently available data at small excess energies by more than two orders of magnitude. It should be noted that both descriptions have been estimated from high energy data, before any of the near-threshold measurements were done.

Close-to-threshold, deviations from the η^4 energy dependence result from the strong s-wave final state interactions in the three two-body pK^+ , $p\Lambda$ and $K^+\Lambda$ subsystems. As the strong interaction in the pK^+ and $K^+\Lambda$ systems appears to be weaker by an order of magnitude as compared to the proton– Λ case [301,302], an accurate description of the energy dependence is expected to be achieved by considering the strong attractive $p\Lambda$ -FSI and repulsive Coulomb corrections in the proton–kaon subsystem. Following the discussion in section 3, a reasonable fit of the energy dependence is obtained by factorizing the reaction amplitude $M_{pp \rightarrow pK^+\Lambda}$ according to eq. (30):

$$\begin{aligned} |M_{pp \rightarrow pK^+\Lambda}|^2 &\approx |M_{FSI}|^2 \cdot |M_0|^2 \cdot F_{ISI} \\ &\approx F_{FSI}(p_{p\Lambda}) F_{Coul}(p_{pK^+}) \cdot |M_0|^2 \cdot F_{ISI} \end{aligned} \quad (44)$$

Here, M_0 is the short-range production amplitude, which is assumed to be constant in the energy range close-to-threshold, M_{FSI} and F_{ISI} describe the elastic interaction in the final state and the reduction due to the pp initial state interaction (see section 3). According to the above discussion the interaction in the three-body final state M_{FSI} is approximated by $F_{Coul}(p_{pK^+})$ and $F_{FSI}(p_{p\Lambda})$, which denote the corrections for the Coulomb interaction and the $p\Lambda$ FSI, depending on the relative momenta p_{pK^+} and $p_{p\Lambda}$ in the proton–kaon and proton– Λ centre-of-mass systems, respectively. The Coulomb correction is

essentially given by the Coulomb penetration factor C^2 of eq. (28)

$$F_{Coul}(p_{pK^+}) = C^2 = \frac{2\pi \eta_{c,pK^+}}{e^{2\pi \eta_{c,pK^+}} - 1}; \quad (45)$$

with the relativistic Coulomb parameter

$$\eta_{c,pK^+} = \alpha \frac{s_{pK^+} - m_p^2 - m_{K^+}^2}{\sqrt{\lambda(s_{pK^+}, m_p^2, m_{K^+}^2)}},$$

where α is the fine structure constant, s_{pK^+} denotes the total energy in the pK^+ subsystem and the triangle function λ is defined in eq. (7).

The modification of the energy dependence close-to-threshold due to the $p\Lambda$ FSI

$$F_{FSI}(p_{p\Lambda}) = \frac{1}{p_{p\Lambda}^2 (1 + \cot^2 \delta_{p\Lambda})} = \frac{1}{p_{p\Lambda}^2 + (-1/\hat{a}_{p\Lambda} + \hat{r}_{p\Lambda} p_{p\Lambda}^2 / 2)^2} \quad (46)$$

is determined by the relative momentum in the $p\Lambda$ centre-of-mass system and the $p\Lambda$ S-wave phase-shift $\delta_{p\Lambda}$, which has been expressed in terms of an effective range expansion in eq. (46), with $\hat{a}_{p\Lambda}$ and $\hat{r}_{p\Lambda}$ as spin-averaged values of $p\Lambda$ scattering length and effective range²². Following the result for the $pp\eta$ final state in eq. (38) in section 3 and using relation (21) from section 2.1, the total cross section for the reaction $pp \rightarrow pK^+\Lambda$ writes as:

$$\sigma(pp \rightarrow pK^+\Lambda) = \frac{F_{ISI} |M_0|^2}{F} \int dV_{ps} \frac{(2\pi \eta_{c,pK^+}) / (e^{2\pi \eta_{c,pK^+}} - 1)}{p_{p\Lambda}^2 + (-1/\hat{a}_{p\Lambda} + \hat{r}_{p\Lambda} p_{p\Lambda}^2 / 2)^2} \quad (47)$$

where F is the flux factor of the colliding nucleons from defined in eq. (9). Using the energy dependence given in eq. (47) an accurate description close-to-threshold is achieved, as shown in figure 14b in section 3 (see also [113]).

Within meson-exchange models, contributions to the reaction dynamics of the elementary strangeness dissociation are expected from direct production (fig. 33a) and from the exchange of nonstrange ($\pi, \eta, \rho, \sigma, \omega$) mesons including N^* resonance excitation and (K, K^*)-exchange in the strange sector (fig. 33b).




Figure 33: Possible production mechanisms for kaon-hyperon production in nucleon-nucleon scattering. a) — Direct kaon emission, b) — Nonstrange and strange meson exchange diagrams, c) — Quark-gluon exchange mechanism. For the latter see the discussion in section 4.1.

²²As the $p\Lambda$ system can be in a spin singlet 1S_0 and spin triplet 3S_1 state, the amplitudes should be added incoherently. However, model calculations [303–305] and experimental data [306,307] suggest that the scattering parameters are rather similar for both spin states (fig. 34).

Pion- and kaon-exchange were suggested as mechanisms of the associated strangeness production in the nucleon-nucleon interaction already by Ferrari [308]. As noted by Laget [310], the contribution from direct production (fig.33a) to the total cross section is small but might be of relevance for differential observables as discussed below.

Modern parameterizations considering π - and K -exchange reproduce the magnitude of the total cross section close-to-threshold within a factor of two or three (grey dashed [297] and grey dotted [298] lines in fig. 32). In refs. [297, 298] the elementary $\pi N \rightarrow KY$ and $KN \rightarrow KN$ transitions are included phenomenologically²³ and both exchange diagrams are found to be important in the case of the reaction $pp \rightarrow pK^+\Lambda$, while Σ production appears to be dominated by π -exchange²⁴. However, interference terms of the amplitudes as well as the effects of final state interactions have been neglected, although the latter have been shown to be of relevance in earlier studies [310].

The same holds for a resonance model (dashed line in fig. 32 [299, 315]) considering π -, η - and ρ -exchange exciting baryon resonances up to 2 GeV/c² coupling to the kaon-hyperon channel (fig. 33b left), i.e. the S_{11} $N^*(1650)$, P_{11} $N^*(1710)$, P_{13} $N^*(1720)$ and P_{33} $\Delta(1920)$ resonances. The fully relativistic model essentially uses parameters fixed by the previous study on π induced strangeness production [311] and experimental branching ratios, interferences of the amplitudes are neglected. Non-resonant exchange diagrams, i.e. especially kaon exchange, are not considered from arguments based on the author's studies on $\pi N \rightarrow KY$ reactions [311]. As a result, π exchange and excitation of the $N^*(1650)$, coupling strongly to the πN channel, is found to give the dominant contribution close-to-threshold and is only exceeded by ρ exchange at excess energies beyond 1 GeV. However, as FSI effects are neglected — according to the authors — the approach is neither expected nor intended to describe the close-to-threshold regime [315].

Within a factor of two, the total cross section data is described by an alternative approach [300] considering π exchange and excitation of the second S_{11} resonance, the $N^*(1650)$ (dotted lines in fig. 32) — in analogy to η production arising from π exchange and excitation of the first S_{11} resonance ($N^*(1535)$). Assuming the $N^*(1535)$ and the $N^*(1650)$ to dominate the ηN and $K^+\Lambda$ systems, respectively, leads to identical forms of the production operators and the spin-angular momentum algebra. Thus, after normalizing the scale of the $pp \rightarrow pK^+\Lambda$ excitation function to low-energy data on the $pp\eta$ final state, the model gives an absolute prediction for Λ production, while the energy dependence close-to-threshold is essentially determined by three-body phase-space modified by the $p\Lambda$ final state interaction. The two corresponding curves shown in figure 32 (dotted lines) differ in the choice of low energy $p\Lambda$ scattering parameters [304–306] determining the proton-hyperon FSI (see eq. (46)).

The situation is summarized by Kaiser [237]: Separating the on-shell S-wave final state interaction and the invariant T matrix at threshold, the information from the close-to-threshold total cross section can be “condensed” into essentially one number, the threshold transition amplitude parametrizing the T matrix, with the energy dependence being accurately reproduced by three-body phase-space and FSI (see also our discussion in section 4.1). In an analysis based on tree-level meson exchange diagrams pointlike ω exchange alone or the total vector ρ^0 , ω and K^* exchange, are found to describe the transition amplitude²⁵. However, also pseudoscalar amplitudes, i.e. pointlike K^+ exchange alone, or — in line with [300, 316] — π exchange followed by an excitation of the S_{11} $N^*(1650)$ resonance when added to

²³For π induced strangeness production, either experimental values [309] (as in the work of Laget [310]) or parametrizations (cross sections determined from the resonance model approach [311] used by Li and Ko [297], and Sibirtsev [298]), respectively, are employed. KN scattering data are available experimentally [312] and in terms of phase-shift analyses [313].

²⁴The relative strength of the K exchange mechanism in Λ and Σ^0 production reflects the smallness of the $KN\Sigma$ coupling constant $g_{KN\Sigma}$ as compared to $g_{KN\Lambda}$ (see e.g. [305, 314])

²⁵The transition amplitudes extracted in [237] based on the approach developed in [34] are related to the production amplitude $|M_0|$ used in this work by simple kinematical factors, i.e. $|M_0|^2 = |\mathcal{T}|^2 \cdot (8 m_p m_B \cdot \lambda(s, m_p^2, m_p^2)) / s$, with m_B denoting the mass of the second baryon $B = p, Y$ in the final state and $|\mathcal{T}|$ as threshold transition amplitude, i.e. $|\mathcal{A}|$ and $|\mathcal{C}|$ for π and η production in [34] and $|\mathcal{K}|/\sqrt{3}$ in [237].

the total pseudoscalar and vector exchange contributions are in good agreement with the experiment. It is concluded [234] that the unpolarized $pp \rightarrow pK^+\Lambda$ total cross section close-to-threshold does not provide enough information to determine the underlying production mechanism.

Instead, further information should be provided in the future by angular distributions and polarization observables, as reported at higher excess energies by the DISTO collaboration [317]: The importance of K exchange for Λ production, which might already be expected from elementary amplitudes ²⁶ has been supported experimentally by data on the normal spin transfer coefficient D_{NN} in exclusive data at an excess energy of $Q = 430$ MeV. D_{NN} is found to be large and the Λ polarization correlated with the beam polarization is oriented opposite to the beam spin for forward Λ production. Based on the model described in [310], calculations by Laget show D_{NN} to discriminate between the π and K exchange mechanism from a different spin coupling at the vertices between the polarized proton in the initial and the hyperon in the final state ²⁷. While from angular momentum and parity conservation π exchange leads to $D_{NN} = +1$ for forward production, kaon emission results in a spin flip ($D_{NN} = -1$) and consequently, the latter is clearly favoured by the experimental result [317] ²⁸. Recently, the negative sign of the spin transfer coefficient, i.e. the dominance of K exchange for Λ production, has been confirmed for two different beam momenta corresponding to excess energies of 319 MeV and 200 MeV, respectively [321]. However, unexpectedly large negative values are also seen for backward production, where the hyperon is preferentially associated with the unpolarized target proton — an observation, which is so far not reproduced within the meson exchange picture.

The first close-to-threshold measurement of the Λ recoil polarization at the COSY-TOF facility has shown a negative polarization for transverse momentum transfers $p_t \geq 0.3$ GeV/c [116] ²⁹, which implies evidence for S- and P-wave interference terms at the respective excess energy of 138 MeV according to [237].

Differential observables, like the occupation in a Dalitz plot, or in projection invariant mass distributions of two-particle subsystems, might also provide important information to identify the relevant production mechanism. A resonant production should influence for example the ΛK^+ invariant mass distribution significantly at excess energies in the range of 100 MeV, as calculated within the resonance model approach of [299, 315] by Sibirtsev et al. [323] ³⁰. First data has already been published [116] and higher statistics distributions at several excess energies will be available from data taken at the COSY-TOF facility in the near future.

Λ production on a neutron target, i.e. via the reactions $pn \rightarrow p(n)K^0(K^+)\Lambda$, might turn out to be sensitive on the relative strength of π , ρ and η exchange, as it has been discussed in case of η production [49, 300].

An important aspect in the theoretical interpretation of data close to the production threshold is the interplay between the type of hyperon-nucleon interaction employed and the result of the model calculation. As already discussed by Laget [310], with experimental observables being significantly influenced by the strong final state interaction at low transverse momenta, exclusive close-to-threshold data should be expected to eventually constrain strongly YN low-energy scattering amplitudes. Within a meson-exchange approach, i.e. considering direct kaon emission as well as π and K exchange (figs. 33a–b) and using an off-shell T matrix description for the hyperon-nucleon interaction the sensitivity on the microscopic hyperon-nucleon potential used has been studied in detail by Kelkar and Jain [325] for Λ production in comparison to the inclusive SATURNE data from [326]. The authors conclude, that — in order to separate different channels contributing at the ΣN threshold — i) exclusive

²⁶The total $\pi p \rightarrow KY$ cross section is in the range of 1 mb compared to 10 mb for K^+p scattering (see fig. 10a).

²⁷For a more general discussion on the interpretation of spin transfer measurements see [318].

²⁸The DISTO results show an opposite sign compared to inclusive D_{NN} high energy ($Q = 16.9$ GeV) data [319], which have been accounted for within quark model calculations [320].

²⁹A negative Λ polarization is also observed for $p_t \geq 0.4$ GeV/c in proton-proton scattering at higher energies [322].

³⁰The calculations on differential distributions have recently been extended to the reaction $pp \rightarrow pK^+\Sigma^0$ in [324].

data is needed (which shall be discussed below) to constrain hyperon–nucleon interaction models and that ii) an improved knowledge on the hyperon–nucleon interaction is prerequisite to further detailed investigations on the relevant production mechanisms. We shall use these conclusions as guideline for the discussions during the next paragraphs.

Due to the short hyperon lifetimes direct YN scattering experiments are difficult to perform and, consequently, low–energy scattering parameters are still rather unknown experimentally. In the case of the Λ –proton system, experimental results have only been available at excess energies above 3.8 MeV [306, 307]. In contrast, a study in close–to–threshold production via the strong final state interaction covers an excess energy range down to threshold, otherwise inaccessible for elastic scattering experiments.




Figure 34: Λp S–wave singlet (open symbols) and triplet (closed symbols) scattering length $a_{p\Lambda}$ and effective range $r_{p\Lambda}$. Experimental data are from Λp elastic scattering (stars [306] and triangles [307]), K^- capture on a deuteron (diamond [327]) and from the FSI approach in threshold production (cross [328], spin–averaged value). Results of YN interaction models are included by squares [303] and circles and double circles (solutions A and B of [304, 305]), respectively. Errors have only been determined for the K^- capture experiment [327].

Assuming the energy dependence of the reaction $pp \rightarrow pK^+\Lambda$ close–to–threshold being determined by eq. (47), i.e. by three–body phase–space modified by the strong $p\Lambda$ FSI and the Coulomb repulsion in the pK^+ subsystem, constraints on the spin–averaged low energy Λp scattering parameters have been extracted in [328] from the differential occupation of Dalitz plot distributions obtained from measurements at the COSY–11 facility [113]. However, in the analysis scattering length and effective range appeared strongly correlated, as inherent to the parametrization used in eqs. (46) and (47). Thus, only a combined fit using the complementary information of low–energy Λp elastic scattering experiments [306, 307] constrained by the Dalitz plot analysis allowed to derive separate spin–averaged values [328]. The result — new spin averaged values of Λp S–wave scattering length and effective range $(\hat{a}_{p\Lambda}, \hat{r}_{p\Lambda}) = (-2.0 \text{ fm}, 1.0 \text{ fm})$ — is shown in figure 34 together with other experimental and theoretical values, where the extracted value of the Λp scattering length is clearly fixed by the old data.

Although the FSI approach to low–energy Λ –proton scattering proves to be a powerful tool, it should be noted that the above values from [328] have been derived assuming a cross section ratio according to the number of possible magnetic spin quantum numbers, i.e. a ratio of three to one between the triplet and singlet Λp final states, which was confirmed in the early bubble chamber experiments.

Consequently, the analysis implies that the underlying reaction mechanism in proton–proton scattering does not favour either the singlet or triplet state, different to the assumed ratio of spin states. Future threshold production experiments employing both polarized beam and target should dispel this concern and allow to extract singlet and triplet hyperon–nucleon scattering parameters separately.

Eventually referring to the first conclusion of Kelkar and Jain [325] cited above, further constraints on the present understanding of close-to-threshold hyperon production have arisen from data on the reaction $pp \rightarrow pK^+\Sigma^0$ taken at the COSY–11 installation [114]: At equal excess energies — up to 13 MeV with respect to the hyperon production thresholds — the total cross sections have been determined for both Λ [113,114] and Σ^0 [114] production. A fit to the data including the hyperon–nucleon FSI is shown in figure 35 and has been performed using the parametrization of the energy dependence developed by Fäldt and Wilkin [110, 120, 300] as given by eq. (35), yet neglecting the weak variation of the cross section due to the variation of the flux factor F . It should be noted that Gasparyan [329] concludes that the proton–hyperon FSI is less important for the ΣN rather than for the ΛN channel. A similar result gives the preliminary analysis of new data taken at COSY–11 for both reaction channels up to excess energies of $Q = 60$ MeV [115].




Figure 35: Total cross sections of the reactions $pp \rightarrow pK^+\Lambda/\Sigma^0$ [113,114]. In comparison, fits of the energy dependence for both hyperon production channels using a pure phase–space calculation (dotted lines) and phase–space modified by the proton–hyperon FSI (solid lines, eq. (35) [110,120,300]) are shown.

The most striking feature of the data is the observed Σ^0 suppression in the energy range close-to-threshold with

$$\mathcal{R}_{\Lambda/\Sigma} = \frac{\sigma(pp \rightarrow pK^+\Lambda)}{\sigma(pp \rightarrow pK^+\Sigma^0)} = 28^{+6}_{-9}, \quad (48)$$

while at excess energies ≥ 300 MeV this ratio is known to be about 2.5 [238]. Considering only π exchange, data on π induced hyperon production via $\pi N \rightarrow K\Lambda(\Sigma^0)$ result in a ratio of $\mathcal{R}_{\Lambda/\Sigma} \approx 0.9$ [330], clearly underestimating the experimental value of relation (48). Kaon exchange essentially relates the ratio $\mathcal{R}_{\Lambda/\Sigma}$ to the ratio of coupling constants squared $g_{\Lambda K}^2/g_{\Sigma K}^2$. Although there is some

uncertainty in the literature on their values [304, 305, 314, 331, 332], a Λ/Σ^0 production ratio of 27 follows from the suitable choice of the SU(6) prediction [232, 333], in good agreement with experiment. However, effects of final state interaction as well as the importance of π exchange especially for Σ^0 production are completely neglected by this simple estimate.

Inclusive K^+ production data taken at the SPES 4 facility at SATURNE in proton–proton scattering at an excess energy of $Q = 252$ MeV with respect to the $pK^+\Lambda$ threshold show enhancements in the invariant mass distribution at the Λp and ΣN thresholds of similar magnitude [326]. However, with only the kaon being detected, it is not clear whether the enhancement at the ΣN threshold is due to Σ production. Qualitatively, a strong $\Sigma N \rightarrow \Lambda p$ final state conversion might account for both the inclusive results from SPES 4 as well as the Σ^0 depletion in the COSY–11 data. In production experiments, evidence for $\Sigma N \rightarrow \Lambda p$ conversion effects is known experimentally from exclusive hyperon data via $K^-d \rightarrow \pi^-\Lambda p$ [327, 334, 335] (for a theoretical interpretation see [336, 337]). In hypernuclear physics, the strong $\Sigma N \rightarrow \Lambda N$ mode appears as the dominating decay channel of Σ –hypernuclei [338], and the ΣN interaction reflects in both shift and broadening of the last X–rays observed from Σ^- atoms [339]. Theoretically, modern hyperon–nucleon interaction models show a significant cusp effect for the ΛN cross section at the ΣN threshold [304, 314].

In exploratory calculations performed within the framework of the Jülich meson exchange model [330], taking into account both π and K exchange diagrams and rigorously including FSI effects in a coupled channel approach, a final state conversion is rather excluded as dominant origin of the observed Σ^0 suppression. While Λ production is found to be dominated by kaon exchange — in agreement with the DISTO results at higher excess energies discussed above — both π and K exchange turn out to contribute to the Σ^0 channel with similar strength. It is concluded by Gasparian et al. [330], that a destructive interference of π and K exchange diagrams might explain the Σ^0 suppression close–to–threshold and a good agreement with the COSY–11 data is obtained after including the overall reduction F_{ISI} in eq. (47) due to the pp interaction in the initial state [19].

An experimental study of Σ production in different isospin configurations should provide a crucial test of the above interpretation: For the reaction $pp \rightarrow nK^+\Sigma^+$ an opposite interference pattern is found, i.e. the $nK^+\Sigma^+$ channel is enhanced for a destructive pattern as compared to the constructive interference of K and π exchange. For the choice of a destructive interference favoured by the $pK^+\Sigma^0$ channel, the corresponding ratio $\sigma(nK^+\Sigma^+)/\sigma(pK^+\Sigma^0) \approx 3$ is in good agreement with high energy data, whereas the alternative choice is rather difficult to reconcile with existing data at higher energies [330]. In contrast, for the reaction $pp \rightarrow pK^0\Sigma^+$ the same interference pattern occurs as for the $pK^+\Sigma^0$ channel, resulting in a ratio $\sigma(pK^0\Sigma^+)/\sigma(pK^+\Sigma^0) \approx 3.3$ close–to–threshold for either choice [340], while a ratio close to unity is found in the literature for higher energies [238].

As stated by the authors of ref. [330], contributions from direct production (fig. 33a) as well as heavy meson exchanges (fig. 33b) have been neglected so far in these calculations but might influence the Λ/Σ^0 production ratio (see also [237, 299, 315]). In fact, employing both a π and K exchange based meson exchange model without any interference of the amplitudes following [298] and the resonance model of [299, 315] the close–to–threshold Λ/Σ^0 production ratio has been studied in [341]. Within a factor of two of the experimental error bars in [114] both models — with parameters fixed at data from excess energies ≥ 1 GeV — are in reasonable agreement with the data, i.e. — according to the authors — the total cross section data close–to–threshold are not sensitive on the details of the model used [341] ³¹.

The latter study has been critically discussed in [342], where the strangeness production is modeled in an effective Lagrangian approach following [107, 316, 343] and strangeness production proceeds via

³¹It should be noted, that unlike [330] π and K exchange amplitudes are concluded by Sibirtsev et al. [341] to contribute to both Λ and Σ^0 production with similar magnitude. As mentioned in [341], an experimental determination of the kaon exchange contribution close–to–threshold — e.g. from polarization observables as in [317] — should be crucial to identify the dominant reaction mechanisms.

π , ρ , ω and σ exchange exciting the nucleon S-wave resonance $N^*(1650)$ and the P-wave resonances $N^*(1710)$ and $N^*(1720)$. For both the Λ and Σ^0 production channel, π exchange followed by an excitation of the $N^*(1650)$ is found to dominate the total cross section close-to-threshold (see also [300]), though an on-shell coupling of the $N^*(1650)$ to the $K^+\Sigma^0$ channel is suppressed with the channel opening at 1686 MeV/c². In contrast, the influence of the $N^*(1650)$ on Σ^0 production and possible interference effects of resonance contributions have been neglected in [341]³². At larger excess energies ($Q \geq 300$ MeV) the $N^*(1710)$ is concluded in the work of Shyam et al. [342] to dominate both reaction channels, which would be expected when considering coupling constants only³³.

It is worth noting, that the OBE calculations performed by Laget [310, 346], although not using the interference of pion and kaon exchange graphs explicitly and just choosing the relative sign to maximize the cross section, not only describe the recent close-to-threshold data on the Λ/Σ^0 ratio within a factor of two, as well as the polarization transfer results of the DISTO experiment [317], but also give an accurate description of the YN invariant mass distributions obtained in the inclusive measurements at SATURNE [326] cited above: In particular, the FSI contribution from the direct $\Lambda p \rightarrow \Lambda p$ amplitude is found to significantly enhance the kaon spectrum just above the Λp threshold, while the strong $\Sigma N \rightarrow \Lambda p$ coupling “induces only a small dip close to the ΣN threshold” in the π and K exchange amplitudes [310]. On the other hand, it is this strong channel coupling and the resulting rapid variation of the 3S_1 ΛN amplitude near the ΣN threshold, which appears responsible for the sharp rise of the invariant mass distribution obtained at SPES 4 at the ΣN threshold via the “kaon direct emission amplitude”, i.e. the coupling of one proton to a K^+Y pair and a following interaction with the second proton (fig. 33a).

³²It has been remarked by Shyam et al. in [342] that the final state interaction employed in [341] is “at variance” with [108] and other approaches in the literature [33, 316, 344].

³³Accordingly, close-to-threshold Σ^0 production in the resonance model of [315] proceeds preferably via π and η exchange, the latter being enhanced for the Σ^0 compared to the Λ channel due to the large branching ratio of the $N^*(1710)$ to the $N\eta$ channel. However, as pointed out in [324] the η exchange contribution is tainted with rather large uncertainties due to different values of the ηNN coupling constant extracted from experimental data (see [345] and section 3).




Figure 36: Λ/Σ^0 production ratio in proton–proton collisions as a function of the excess energy. Experimental data within the shaded area, which corresponds to the range described by relation (48), are taken from [114], data at higher excess energies from [115]. Calculations [347] are performed within the Jülich meson exchange model, assuming a destructive interference of K and π exchange and employing the microscopic YN interaction models Nijmegen NSC89 (solid line [314]) and the new Jülich model (dotted line [348]), respectively.

Recently, measurements of the Λ/Σ^0 production ratio in proton–proton collisions have been extended up to excess energies of $Q = 60$ MeV at the COSY–11 installation [115], in order to study the transition between the low–energy enhancement of the ratio reported in [114] and data at excess energies larger than 300 MeV. In comparison to the experimental data, in figure 36 calculations are included obtained within the approach of Gasparian et al. [330] assuming a destructive interference of π and K exchange with different choices of the microscopic hyperon nucleon model [347].

As it was already emphasized in [330], the result depends on the details — especially the off–shell properties — of the microscopic hyperon–nucleon interaction employed, although the actual choice does not alter the general result in [330] of only the destructive interference of K and π exchange explaining the experimentally observed suppression of the Σ^0 signal close–to–threshold. At the present stage of theoretical investigations, both the good agreement found for Jülich model A [304] with the close–to–threshold result for $\mathcal{R}_{\Lambda/\Sigma}$ in relation 48 and for the Nijmegen model (solid line in fig. 36 [314]) with the energy dependence of the cross section ratio in figure 36 should rather be regarded as accidental. In the latter case an $SU(2)$ breaking in the 3S_1 ΣN channel had to be introduced [314]. Consequently, the relation between the $\Sigma^0 p$ amplitude with the $\Sigma^+ p$ and $\Sigma^- p$ channels becomes ambiguous. Only one of the choices leads to the good agreement with the data shown in figure 36 (solid line), whereas the other one results in a completely different prediction [349].

Calculations using the new Jülich model (dotted line in figure 36 [348]) do not reproduce the tendency of the experimental data. It is suggested in [347] that neglecting the energy dependence of the elementary amplitudes and assuming S –waves in the final state might no longer be justified beyond excess energies of 20 MeV.

Conclusions from figure 36, as well as considerations by Kelkar and Jain [325] for the $pp \rightarrow pK^+\Lambda$ reaction clearly demonstrate, that — once the reaction mechanism for close–to–threshold hyperon production is understood — exclusive hyperon production data should provide a strong constraint on the details of hyperon–nucleon interaction models.

Finally, in future high resolution hyperon production experiments close to threshold may give further

information on the possible existence of low-lying strange dibaryons. Narrow, stable six-quark objects D_s and D_t with a $q^4 \otimes q^2$ cluster substructure and strangeness $S = -1$ coupling to P-wave nucleon-hyperon systems have been postulated from an extended MIT bag-model [350] below and above the ΣN thresholds with widths in the range of 10 keV to 10 MeV. Experimental signatures consistent with a strange dibaryon hypothesis have been found both in strangeness transfer reactions $K^- d \rightarrow \pi^- Y N$ [327, 351, 352] and in the inclusive $pp \rightarrow K^+ X$ data taken at SATURNE [326] at excess energies of 175 and 306 MeV with respect to the $\Sigma^0 p$ threshold. However, the statistical accuracy in these experiments turned out to be too small for unequivocal conclusions. For a high resolution study of the inclusive $pp \rightarrow K^+ X$ reaction channel new experimental efforts are on their way at the BIG KARL spectrometer at COSY, with a resolution improved by a factor of five and a higher statistical accuracy compared to the SATURNE measurements [353].

5 Meson pair production

5.1 Double pion production in NN scattering

Single and double pion production in elementary πN , γN and NN reactions is an important source to gain information on nucleon-nucleon, nucleon-pion and pion-pion interactions. Apart from these fundamental aspects, the pion production on nucleons also enables the study of the properties of nucleonic resonances, which might be excited in the reaction processes. The reaction $\gamma N \rightarrow N\pi\pi$ for instance has been found to be influenced by the formation of the $N^*(1520)$ resonance [354] with a subsequent decay into $\Delta\pi$. Therefore, studies on this reaction channel allow to gain information about the $N^*(1520) \rightarrow \Delta(1232)\pi$ decay amplitudes. Additionally, contributions of the $N^*(1440)$ resonance, followed by its decay $N^*(1440) \rightarrow N(\pi\pi)_{S\text{-wave}}^{T=0}$, play a non-negligible role in the near-threshold production of pion-pairs via the reaction channels $\gamma N \rightarrow N\pi\pi$ and $\pi N \rightarrow N\pi\pi$ [355].

In case of the pion-pair production in the nucleon-nucleon scattering similar contributions of nucleonic resonances are under discussion. Recently, detailed calculations on the two-pion production in nucleon-nucleon interactions have been performed [356], considering non-resonant terms as well as contributions from nucleonic resonances with subsequent decays into a nucleon and pions. One of the main results of these studies is the prediction of a dominant effect of the $N^*(1440)$ Roper resonance, followed by its decay $N^*(1440) \rightarrow N(\pi\pi)_{S\text{-wave}}^{T=0}$, in the region of low excess energies. According to this, the absolute scale of the low energy total cross section of reaction channels, where the pions can be in an isospin zero state, e.g. $pp \rightarrow pp\pi^+\pi^-$ and $pp \rightarrow pp\pi^0\pi^0$, should be determined by the excitation of this resonance. In this region of excess energies contributions of non-resonant terms are discussed to be negligible for the description of the excitation function of both these reaction channels. However, at higher energies ($Q > 200$ MeV) the excitation of Δ resonances has been evaluated to dominate the total cross sections.

A different situation is given in the case of the $pp \rightarrow pn\pi^+\pi^0$ reaction channel. Here contributions of the Roper resonance with the subsequent decay into $N(\pi\pi)_{S\text{-wave}}^{T=0}$ are forbidden by conservation laws and contributions of remaining N^* production diagrams are suppressed at low excess energies. Since the non-resonant terms are also negligible in this case, the dominating processes are given by Δ resonance excitations.

It is worth noting that the discussed model calculations yield predictions for the excitation functions for different two-pion production reactions from threshold up to high excess energies ($Q \approx 300$ MeV) and all are based on the same Feynman diagrams. However, especially in the close-to-threshold region the data situation for most double pion reaction channels is far from being optimal. Recently, three isospin-independent reaction channels on the two-pion production in proton-proton interactions, $pp \rightarrow pp\pi^+\pi^-$, $pp \rightarrow pp\pi^0\pi^0$ and $pp \rightarrow pn\pi^+\pi^0$, have been studied in high statistics precision measurements at the

PROMICE/WASA facility at CELSIUS and resulted in the first total and differential cross sections in the near-threshold region [357, 358].

In fig. 37a the total cross sections for the $\pi^+\pi^-$ production in proton-proton scattering are presented up to an excess energy of $Q \approx 300$ MeV. The solid and the dashed lines represent the above discussed predictions of ref. [356] for this reaction channel and correspond to two possible solutions in the applied model. Obviously these calculations are in good agreement with the experimental data both close-to-threshold and in the region of higher excess energies, while the model seems to underestimate the data at intermediate energies. It should be noted that the data points from Dakhno et al. [362] have been obtained using a deuterium filled bubble chamber and selecting events from the quasi-free two-pion production $pp(n) \rightarrow pp(n)\pi^+\pi^-$. However, in the analysis of the data the Fermi motion of the nucleons in the deuteron has been neglected. In later calculations an adequate consideration of the deuteron wave function has been found to shift the effective excess energy by ~ 20 MeV towards higher energies [365], corresponding to a shift of the data towards the region expected by the discussed model calculation.




Figure 37: Total cross section data for the reactions:

- a) $pp \rightarrow pp\pi^+\pi^-$,
- b) $pp \rightarrow pp\pi^0\pi^0$ and
- c) $pp \rightarrow pn\pi^+\pi^0$.

The solid and the dashed curves represent model calculations from [356].

The data were taken from [357–364].

A version of the article including a figure of better quality can be found at the COSY-11 homepage:

http://ikpe1101.ikp.kfa-juelich.de/cosy-11/pub/List_of_Publications.html#papers

Figure 37b displays the experimental situation for the reaction channel $pp \rightarrow pp\pi^0\pi^0$. The new data from PROMICE/WASA disfavour the solution of the model calculation indicated by the solid line but are in very good agreement with the prediction shown by the dashed line. However, the high energy data from ref. [363] are in disagreement with both curves and are clearly overestimated.

Finally, the close-to-threshold total cross section data of the reaction channel $pp \rightarrow pn\pi^+\pi^0$ are presented in fig. 37c.

Different to the previous situation now the data is underestimated by both solutions of the model calculations, especially in the near-threshold region. Since now a production via the $N^*(1440) \rightarrow N(\pi\pi)_{S\text{-wave}}^{T=0}$ is forbidden, this observation might indicate contributions from processes like $NN \rightarrow NN^* \rightarrow N\Delta\pi \rightarrow NN\pi\pi$ larger than predicted by theory.

It should be mentioned that contributions of higher partial waves and final state interactions have been neglected in the model calculations, indicated by the presented curves. However, the effect of the pp FSI has been investigated and is reported to increase the close-to-threshold cross sections by nearly one order of magnitude, while at $Q \sim 120$ MeV this effect is reduced to an increase of a factor of ~ 2 [366]. According to this, considering the interaction of the final state baryons is expected to reduce the apparent discrepancy between the $\pi^+\pi^0$ total cross section data and the corresponding model calculations. Surprisingly, in case of the $\pi^+\pi^-$ and $\pi^0\pi^0$ pair production, the near-threshold data are described reasonable by these calculations neglecting the pp FSI.

Furthermore, signals for the contribution of higher partial waves are reported for the near-threshold data [357], which also have been neglected in the discussed model calculations. This observation might be interpreted as a signal of a heavy meson exchange (σ , ρ) between the interacting protons.

5.2 Double pion production in pd scattering

It is well known and cited in the literature as the ABC-anomaly [367, 368] that the missing mass enhancement at around 310 MeV with a width of 50 MeV observed in the inclusive measurements of the $pd \rightarrow {}^3\text{He} X^0$ reaction at a proton beam energy of $T_p \geq 745$ MeV (corresponding to an excess energy of $Q \geq 190$ MeV for the $\pi\pi$ production) is not excited in the ${}^3\text{H} X^+$ final state. Detailed investigations confirm that this observation is to be associated with the isospin-zero s-wave $\pi\pi$ double-pion state but additionally stress that the missing mass peak and its width vary with the beam momentum [369]. Further, since the isoscalar $\pi\pi$ scattering length is small [370] the observed anomaly is supposed to be of kinematical origin and associated with the intermediate excitation of two Δ resonances [371].




Figure 38: Differential cross sections for the $pd \rightarrow {}^3\text{He} \pi^+ \pi^-$ reaction at $T_p = 546$ MeV as a function of four kinematical variables. Dashed curves represent phase-space predictions normalized to the data whereas the solid curves are calculations assuming the $\pi^+ \pi^-$ system to emerge in a relative p-wave, see [372].

Recently the MOMO (**M**onitor-**o**f-**M**esonic-**O**bservables) collaboration at COSY investigated the $\pi\pi$ production closer to threshold in an exclusive measurement of the $pd \rightarrow {}^3\text{He} \pi^+ \pi^-$ reaction at a beam energy of $T_p = 546$ MeV (excess energy $Q = 70$ MeV for the $\pi\pi$ production), where the ABC enhancement is expected to show up near the centre of the available $\pi\pi$ -invariant mass distribution [372].

In figure 38a–d differential cross sections in terms of four kinematical variables are presented:

- a) The $\pi^+ \pi^-$ excitation energy $T_{\pi\pi} = m_{\pi\pi} - 2m_\pi$, where $m_{\pi\pi}$ is the two-pion invariant mass.
- b) The π - ${}^3\text{He}$ excitation energy.
- c) The angle (θ_π) between the proton and one of the π 's in the overall centre-of-mass (c.m.) system.
- d) The angle ($\theta_{\pi\pi p}$) between the relative $\pi^+ \pi^-$ momentum and the beam axis in the overall c.m. system.

The $T_{\pi\pi}$ distribution at this near-threshold data is pushed closer to the maximum values of excitation energy and is in contrast to the original ABC result [367,368] which showed an enhancement over phase-space at $T_{\pi\pi} \approx 30$ MeV. The distribution of the π - ${}^3\text{He}$ excitation energy is rather consistent with the phase-space prediction. The significant anisotropy of θ_π confirms the importance of higher partial waves involved in the reaction process. The MOMO collaboration concludes [372] that the agreement between the $T_{\pi\pi}$ data of fig. 38a with the solid line (phase-space multiplied by $T_{\pi\pi}$) and the fair linearity of $\sin^2 \theta_{\pi\pi p}$ indicates that the $\pi^+ \pi^-$ system is essentially produced with an internal angular momentum $l = 1$.

On the other hand, as outlined in section 5.1, the Valencia group [356,373] developed a model for the two pion production resulting in a dynamical origin for the small invariant masses of the low energy

part, where the two pions can be in an isospin zero state. Here the reaction dynamics is dominated by the intermediate excitation of the $N^*(1440)$ Roper resonance followed by the $N^* \rightarrow \Delta\pi \rightarrow N(\pi\pi)_{s-wave}^{T=0}$ decay.

The experimental situation seems to be the following: s-wave $\pi\pi$ production has been observed in the $pd \rightarrow {}^3He \pi^+\pi^-$ reaction very close-to-threshold [374] and at high excess energies ($Q \approx 200$ MeV) [367, 368] whereas at an intermediate excess energy range a p-wave $\pi^+\pi^-$ production process dominates. Similar results have been obtained in $np \rightarrow dX$ [375,376] and in the $\pi^+d \rightarrow \pi^+\pi^+nn$ ($\pi^+\pi^-pp$) [377,378] reactions.

5.3 Double K-meson production in pd scattering

After the remarkable results of p-wave production in the $\pi^+\pi^-$ system the MOMO collaboration continued equivalent measurements for the $pd \rightarrow {}^3He K^+K^-$ reaction at excess energies of $Q = 56$ MeV, 40 MeV and 35 MeV.

The K^+K^- invariant mass spectra in units of the K^+K^- relative energy $T_{K^+K^-}$ are shown in figure 39. A clear signal of the ϕ -meson production is seen above the continuous non-resonant K^+K^- spectrum. The dashed lines in all three parts of the figure correspond to simple phase-space calculations, where the Q -dependences of the cross sections ($\sigma_{K^+K^-}$ and σ_ϕ) were taken to scale as $\sigma_{K^+K^-} \propto Q^2$ for the three-body and $\sigma_\phi \propto Q^{1/2}$ for the two-body final state, respectively. The relative angular momentum between the two kaons was assumed to be $l = 0$. No significant deviation of the data from these summed phase-space distributions is observed and thus it is interesting to note, that no indication of p-wave in the K^+K^- system is present in the mass spectra and angular distributions, in contrast to the above discussed data for the two pion production via the reaction $pd \rightarrow {}^3He \pi^+\pi^-$.




Figure 39: K^+K^- invariant mass spectra from the reaction $pd \rightarrow {}^3\text{He} K^+K^-$ at $Q = 56$ MeV, 40 MeV and 35 MeV above threshold plotted in units of the K^+K^- relative energy $T_{K^+K^-}$, see [379].

5.4 Strangeness dissociation into $K\bar{K}$ -pairs

Measurements on the $K\bar{K}$ meson pair production are of considerable interest in the context of the continuing discussion on the nature of the scalar resonances $f_0(980)$ and $a_0(980)$, as already discussed in section 4.3.

Available close-to-threshold data [77, 78] on the elementary antikaon production channel in proton–proton scattering, $pp \rightarrow ppK^+K^-$, are shown in figure 40. In comparison, parametrizations of the energy dependence of the total cross section based on a four–body phase–space behaviour and for a resonant production via the f_0 are included. Obviously, within the experimental error bars and neglecting the possible influence of higher partial waves ³⁴, the data is consistent with both a non–resonant and resonant production, as already concluded from figure 31 in section 4.3, i.e. from the K^+K^- invariant mass distribution obtained at an excess energy of $Q = 17$ MeV [273].

³⁴Preliminary data by the DISTO collaboration [380] indicate that K^+K^- production at an excess energy of $Q = 114$ MeV is still consistent with isotropic emission.




Figure 40: Total cross sections for the free K^+K^- pair production in proton-proton collisions [77, 78]. The solid line indicates calculations on the basis of the non-resonant four-body phase-space expectations fitted to the data points. The dashed lines correspond to three-body phase-space for the resonant K^+K^- production via the f_0 , normalized to the COSY-11 data point). The pp FSI and the Coulomb interaction have been taken into account. The effect of the large uncertainty about the width of the $f_0(980)$ resonance ($\Gamma = 40 - 100$ MeV [134]) is indicated by the dashed area.

A version of the article including a figure of better quality can be found at the COSY-11 homepage:

http://ikpe1101.ikp.kfa-juelich.de/cosy-11/pub/List_of_Publications.html#papers

Furthermore, exclusive K^- production data are of special interest with respect to subthreshold kaon production experiments in nucleus-nucleus interactions, which are expected to probe the antikaon properties at high baryon density. Recent inclusive subthreshold measurements [291–293] resulted in comparable K^+ and K^- yields at the same energy per nucleon below the production thresholds for the elementary reactions $pp \rightarrow K^\pm X$, which has been discussed as an indication for a possible restoration of chiral symmetry (as reviewed in [290]).

Available inclusive elementary K^+ and K^- total cross sections in nucleon-nucleon interactions are shown in figure 41. In contrast to the results obtained in subthreshold measurements, in proton-proton scattering the inclusive K^+ cross section exceeds K^- production by more than an order of magnitude in the near threshold region. To explain this observation, different models [74, 286–289, 381, 382] consider kaons (K^+ , K^0) and antikaons (K^- , $\bar{K}^0$) to be subject of repulsive and attractive forces within the nuclear medium, respectively. Recent theoretical studies using a chiral unitary approach to describe the K^-N S-wave interaction allow for a self-consistent microscopic implementation of medium effects [286, 287]. The K^- self-energies obtained are consistent with kaonic atom data, as shown in [383].




Figure 41: World data on the inclusive total K^+ and K^- production cross sections in proton–proton collisions at excess energies below $Q = 2$ GeV. The data are taken from [77, 113–118, 238, 317, 384].

6 Meson production on light nuclei

Near-threshold meson production experiments in nucleon–nucleon interactions might help to obtain fundamental information on the elementary production processes and final state interactions. Extending those measurements from the binary nucleon scattering to reactions with three or four participating nucleons, i.e. meson production in the proton–deuteron or deuteron–deuteron scattering, allows to study the role of further participating nucleons. This facilitates a gain in information on possible multi-step production processes, which are naturally absent in the nucleon–nucleon case.

6.1 Dynamics in three and four nucleon systems

Of considerable interest is the production of η mesons in the $pd \rightarrow {}^3\text{He} \eta$ reaction. Studied at the SATURNE accelerator in the near threshold region [385, 386] this channel is reported to expose remarkable features. In spite of the much higher momentum transfer, the observed threshold amplitude f_η for the η production, defined by the unpolarized centre-of-mass cross section

$$\frac{d\sigma}{d\Omega}(pd \rightarrow {}^3\text{He} \eta) = \frac{p_\eta^*}{p_d^*} |f_\eta(pd \rightarrow {}^3\text{He} \eta)|^2, \quad (49)$$

was found to be comparable to that for the $pd \rightarrow {}^3\text{He} \pi^0$ reaction at its respective threshold [385]. In addition, while being consistent with s-wave production, $|f_\eta(p_\eta^*)|^2$ decreases by a factor of three from threshold up to a centre-of-mass momentum of emitted η mesons of $p_\eta^* = 70$ MeV/c ($\hat{=}$ $Q \sim 7$ MeV).

This observed strong decrease in the near-threshold region can be addressed to a strong η ${}^3\text{He}$ final state interaction and is therefore of special interest in view of the existence of η –mesic nuclei. Although being the topic of several theoretical investigations [138, 140, 387–391], the possibility for the formation of quasi-bound ${}^3\text{He} \eta$ states is still an open question.




Figure 42: Total cross section σ and square of the production amplitude f for the $pd \rightarrow {}^3\text{He} \eta$ reaction. The lines are explained in the text. The data are from references [386, 392, 393] and show only the statistical uncertainties.

A version of the article including a figure of better quality can be found at the COSY-11 homepage:
http://ikpe1101.ikp.kfa-juelich.de/cosy-11/pub/List_of_Publications.html#papers

The experimental situation for the $pd \rightarrow {}^3\text{He} \eta$ reaction is presented in fig. 42, displaying total cross section data from threshold up to an excess energy of $Q = 100$ MeV.

The SATURNE data [386] (circles) expose an energy dependence, which clearly differs from pure phase-space expectations: According to naive two-body phase-space considerations, neglecting higher partial waves, an excitation function according to $\sigma \propto Q^{1/2}$ is expected. One promising ansatz to describe the reaction process was given by Kilian and Nann [394], taking into account both nucleons from the target and considering double-scattering diagrams. A sketch of such a double-scattering diagram is illustrated in the left part of fig. 43.




Figure 43: Production diagrams for the $pd \rightarrow {}^3\text{He} \eta$ reaction.

Calculations by Fäldt and Wilkin [395] are in line with this approach and describe the $pd \rightarrow {}^3\text{He} \eta$ reaction as an initial $pp \rightarrow d\pi^+$ ($pn \rightarrow d\pi^0$) production followed by a final reaction according to $\pi^+n \rightarrow \eta p$ ($\pi^0p \rightarrow \eta p$). These investigations lead to a prediction of the spin averaged threshold

amplitude squared of $|f(p_\eta^* = 0)|^2 = 0.69 \mu\text{b/sr}$, underestimating the measured results by a factor of 2–3. Since none of the considered sub-reactions exposes a significant variation of the cross section in the relevant energy region, the observed energy dependence of the production amplitude f_η has been interpreted to be a signal of a strong final state interaction of the η -nucleus system.

Introducing the ${}^3\text{He} \eta$ final state interaction by an s-wave scattering length formula

$$f(p_\eta) = \frac{f_B}{1 - i a_{{}^3\text{He} \eta} p_\eta^*}, \quad (50)$$

with f_B as the threshold amplitude, the best description of the observed energy dependence of the SATURNE total cross section data is achieved using a scattering length of $a_{{}^3\text{He} \eta} = \pm(3.6 \pm 0.5) \text{ fm} + i(2.0 \pm 0.9) \text{ fm}$, as can be seen in fig. 42 (solid line) with the absolute scale of the curve being scaled to fit the data. Such a large scattering length might be associated with the existence of a quasi-bound ${}^3\text{He} \eta$ state.

In the right part of fig. 42 the extracted production amplitude squared $|f(p_\eta^*)|^2$ is presented as function of the centre-of-mass momentum of the ejectiles. It should be noted that in the close vicinity of the production threshold and with a given cross section, the production amplitude squared $|f(p_\eta^*)|^2$ depends crucially on the momentum p_η^* and, therefore, on the exact masses of the ejectiles. Here a mass of the ${}^3\text{He}$ -nucleus of $m_{{}^3\text{He}} = 2.808392 \text{ GeV}/c^2$ [396] and a meson mass of $m_\eta = 0.54730 \text{ GeV}/c^2$ [134] have been applied. While a constant amplitude would imply a behaviour according to pure s-wave phase-space considerations, the observed shape exposes a strong momentum dependence and can be reproduced well by the discussed formalism up to excess energies of $Q \approx 7 \text{ MeV}$, corresponding to $p_\eta^* \approx 0.35 \text{ fm}^{-1}$.

However, recent measurements at higher excess energies from the GEM collaboration at COSY [392] and from the WASA/PROMICE collaboration at CELSIUS [393] clearly deviate from the predicted energy dependence of the cross section (fig. 42). Moreover, different to the SATURNE results, the centre-of-mass angular distribution of the emitted η mesons strongly differs from an isotropic emission and exposes a peaking in forward direction. Motivated by results on the η meson production in γp and pp interactions, another approach to describe the production process has been evaluated [392] where the η meson creation is described by the excitation of the $S_{11} N^*(1535)$ resonance as shown in figure 43 (right). Neglecting effects of other resonances, the production amplitude can be expressed by a Breit-Wigner distribution [397]:

$$|f(E)|^2 \sim \left| \frac{E_R \Gamma_R}{(E_R^2 - E^2) - i E_R \Gamma(E)} \right|^2, \quad (51)$$

with E_R as the mass of the S_{11} resonance and Γ_R as the corresponding width. Since the production threshold is located in the vicinity of the resonance mass E_R , the energy dependence of the width $\Gamma(E)$ has to be considered:

$$\Gamma(E) = \Gamma_R \left(b_\eta \frac{p_\eta^*}{p_{\eta R}^*} + b_\pi \frac{p_\pi^*}{p_{\pi R}^*} + b_{\pi\pi} \right). \quad (52)$$

The parameters b_i indicate the branching ratios for different decay modes of the resonance while the centre-of-mass momenta of the η and π mesons are represented by p_i^* . A corresponding calculation, scaled to fit the GEM data point, is given in figure 42 by the dash-dotted line. Obviously, the prediction of this approach fails to describe the shape of the SATURNE data. This observation might indicate an influence of a strong ${}^3\text{He} \eta$ final state interaction only in the close vicinity of the production threshold.

Alternatively, this effect might be caused by a transition between different production processes.

Concluding, the $pd \rightarrow {}^3\text{He}\eta$ reaction appears to be an interesting and important channel to study both the meson–nucleus final state interaction as well as reaction processes involving three nucleons. However, the body of data is far from being optimal, especially in the near–threshold region. New data from PROMICE/WASA and COSY–11 is currently under evaluation and will improve the experimental situation.

6.2 Quasi–bound states

From the theoretical point of view the existence of mesic nuclei or quasi–bound meson–nucleus systems is not excluded, however, up to now a compelling experimental proof for the formation of such a state is still missing. It is obvious that the possibility to create such states crucially depends on both the properties of the meson and the nucleus as well as on the sign and the strength of the meson–nucleus interaction. Since the π –nucleon and K^+ –nucleon interaction have been found to be repulsive, it is unlikely that they will form quasi–bound states. Contrary, there are evidences for an attractive K^- –nucleon interaction, however, corresponding experiments would suffer from the low production cross sections for K^- mesons. Furthermore, the Coulomb interaction is expected to screen corresponding physical observables and might lead to the formation of mesonic atoms bound by the Coulomb interactions. Different to this, the η meson is uncharged and the observation of an attractive η –nucleon interaction led to speculations concerning the existence of η –nuclear quasi–bound states. In the absence of η –meson beams such states bound by the strong interaction would offer a new possibility to study the η –nucleon interaction since the meson would be trapped for a relatively large time in nuclear medium. Predicted by Haider and Liu [387, 388], detailed studies on the formation of such states have been the topic of recent investigations [138, 140, 389–391, 398], resulting in different lower limits of the atomic number A for which bound states should exist. While e.g. in [387, 388] the lower mass number is estimated in the framework of an optical model approach to be $A = 12$, there are considerations that even for nuclei of lower masses ($d, t, {}^3\text{He}, {}^4\text{He}$) bound systems might exist [138, 140], see also section 3.

Experimental evidences for bound states have been found in the near–threshold production of η mesons in the reaction channel $pd \rightarrow {}^3\text{He}\eta$. The unexpected large production amplitude as well as its rapid decrease with increasing energy (figure 42) is attributed to a strong s–wave final state interaction associated with a large η – ${}^3\text{He}$ scattering length [140].

Such a large scattering length (see fit in figure 42) might imply the existence of a quasi–bound state. Assuming the ${}^3\text{He}\eta$ to form a quasi–bound state, it is therefore interesting to compare $pd \rightarrow {}^3\text{He}\eta$ production data with the ones of the reaction $dd \rightarrow {}^4\text{He}\eta$, where according to the higher mass number, the formation of a bound state should be even more probable. A comparison of corresponding near–threshold data is presented in figure 44.

A fit to the data using the scattering length formula (50) provides information on the scale of both the real and imaginary part of the scattering length of the meson–nucleus system but is insensitive to the sign of the real part of $a_{\text{He}\eta}$. However, information on this sign can be obtained using a lowest–order optical potential [140]:

$$2 m_R V_{\text{opt}}(r) = -4\pi A \rho(r) a_{\eta N} \quad (53)$$

with the η –nucleon reduced mass m_R , the mass number of the nucleus A and the η –nucleon scattering length $a_{\eta N} \approx (0.52 + i 0.25)$ fm. Using Gaussian nuclear densities with rms radii of $\rho(r) = 1.63$ fm

(1.9 fm) for the $\eta^3\text{He}$ ($\eta^4\text{He}$) system, meson–nucleus scattering lengths of

$$a(^3\text{He}\eta) \sim (-2.3 + i 3.2) \text{ fm} \quad (54)$$

$$a(^4\text{He}\eta) \sim (-2.2 + i 1.1) \text{ fm} \quad (55)$$

with negative real parts for both systems have been obtained.




Figure 44: Averaged squared production amplitudes of the reactions $pd \rightarrow ^3\text{He}\eta$ and $dd \rightarrow ^4\text{He}\eta$ as function of the centre-of-mass η momentum. The curves are discussed in the text. The data are taken from [386, 399, 400].

Fits to existing close-to-threshold data for both reaction channels using these scattering lengths and eq. (50) are presented in figure 44. Obviously, the data are reproduced well by this formalism. These results imply the existence of quasi-bound meson–nucleus states in the systems $\eta^3\text{He}$ and $\eta^4\text{He}$, however, unambiguous evidence is still missing. Therefore, detailed investigations on both reaction channels as well as on reactions using heavier targets are highly recommended.

6.3 Test of invariances

Conservation laws restrict the quantum numbers for the production of mesons in the nucleon–nucleon interaction. Here especially the pion production is of considerable interest since it exhausts all inelasticity up to about 2 GeV/c beam momentum in the nucleon–nucleon collision.

As outlined in references [4, 5, 401, 402] the conservation of isospin allows the total cross section for pion production to be expressed as an incoherent sum of three independent partial cross sections given by: σ_{11} , σ_{10} and σ_{01} , where the first (second) index indicates the isospin of the initial state (final state) two–nucleon system and σ_{00} is ruled out due to isospin conservation arguments.

The three cross sections can be determined via the reactions:

pure σ_{11} channels:	$\sigma(pp \rightarrow pp\pi^0)$,	$\sigma(nn \rightarrow nn\pi^0)^*$	
pure σ_{10} channels:	$\sigma(pp \rightarrow d\pi^+)$,	$\sigma(np \rightarrow d\pi^0)$,	$\sigma(nn \rightarrow d\pi^-)^*$
mixed $\sigma_{10} + \sigma_{11}$ channels:	$\sigma(pp \rightarrow pn\pi^+)$,	$\sigma(nn \rightarrow np\pi^-)^*$	
mixed $\sigma_{10} + \sigma_{01}$ channels:	$\sigma(np \rightarrow pn\pi^0)$,		
mixed $\sigma_{11} + \sigma_{01}$ channels:	$\sigma(np \rightarrow pp\pi^-)$,	$\sigma(np \rightarrow nn\pi^+)$.	

Charge symmetry relates those reactions denoted by a * to their mirror systems given as the first one in the same row.

The two reactions $pd \rightarrow {}^3H \pi^+$ and $pd \rightarrow {}^3He \pi^0$, which permit the study of isospin breaking effects have been studied systematically [403]. Figure 45 compares the differential cross sections for these two reactions at two proton momenta close to the production threshold. At these momenta no significant differences in the differential cross sections were observed. In fact, interpolating the cross sections to the same four-momentum-transfer yields an expected average ratio $R = \sigma(pd \rightarrow {}^3H \pi^+)/\sigma(pd \rightarrow {}^3He \pi^0) \approx 2.0$ within the experimental uncertainties. However, the COSY-GEM collaboration seems to have indications that a clear effect of isospin breaking is observed at a momentum of about 1.57 GeV/c, close to the η -production threshold [404].




Figure 45: Differential cross sections for the two isospin symmetric reactions $pd \rightarrow {}^3He \pi^0$ and $pd \rightarrow {}^3H \pi^+$ at beam momenta of 800 MeV/c and 750 MeV/c, respectively. Data are from the GEM collaboration [403], the error bars are smaller than the symbols.

As pointed out in reference [405, 406], in context of QCD isospin and charge symmetry are broken due to both i) the mass difference between up and down quarks and ii) the electromagnetic interaction. The observation of mixing between mesons belonging to the same $SU(3)$ multiplets offers a unique possibility to study charge symmetry breaking effects. As suggested by Wilkin [407] a possible isospin symmetry breaking can manifest itself e.g. via external π^0 - η mixing. Searching for such effects the authors of reference [405] are investigating the cross section ratio for the $pd \rightarrow {}^3He \pi^0$ and $pd \rightarrow {}^3H \pi^+$ reactions in particular for large relative pion-proton angles and at beam energies of the corresponding η production threshold. The idea is based on the assumption that — besides the direct reaction — an intermediately formed $\tilde{\eta}$ meson undergoes a transition to the final π^0 -meson for the $pd \rightarrow {}^3He \pi^0$ reaction ($pd \rightarrow {}^3He \tilde{\eta} \rightarrow {}^3He \pi^0$) with a yield governed by the strength of the mixing angle while such an intermediate state can not occur for the $pd \rightarrow {}^3H \pi^+$ case. The authors [405] predict angular

dependent cross section ratios as: $R_{\theta_{p\pi}=180^\circ} = 2.4$ and $R_{\theta_{p\pi}=0^\circ} = 2.03$. The predicted symmetry breaking magnitude appears to be in reach of the experiments planned by the GEM collaboration at COSY.

Another direct evidence for the isospin invariance breaking would be the observation of the non-zero yield for the isospin forbidden $dd \rightarrow {}^4He \pi^0$ reaction. The upgrade of the ANKE detection system at COSY will allow to search for the effect [408].

7 Experimental facilities

Studies of meson production at threshold require crucially high quality and precise quantitative knowledge of the accelerator beam quality especially due to the strong cross section dependence on the excess energy. The rapid growth of the meson production cross section in the vicinity of the kinematical threshold is evidently seen in figure 2. It has been shown in the previous sections that new precise data are available for different reaction channels. Here we present a very brief overview of technical features of such experimental facilities as: accelerators, targets and detector arrangements where only a representative selection can be given.

7.1 Accelerators for hadronic physics at medium energies

Initiated by the rather high costs for experimenting with antiprotons and aiming in a most efficient usage, storage rings have been developed. The possibility of controlling the emittance of such beams by electron as well as stochastic cooling, combined with some unique benefits of using internal targets lead to the construction of medium energy cooler rings as the low energy antiproton ring (LEAR) which might be regarded as the father for many facilities built.




Figure 46: Accelerators for medium energy physics.

The maximum available beam momentum limits the producible mass. In figure 46 a comparison is presented between proton and electron accelerators in the range of producing in the proton-proton and

γ -proton collision mesons up to the ϕ -meson at 1.02 GeV/c². This self-explaining figure demonstrates e.g. that the Uppsala accelerator CELSIUS can reach beam momenta to produce the η -meson in the proton-proton scattering which is about the same mass production range as that of MAMI via the γ -proton interaction.

Complementary to the investigations with charged hadronic beams the high precision secondary neutron beam of the TRIUMF facility [46] permits the close-to-threshold meson production in neutron-proton collisions.

In Table 7.1 some accelerators used for hadron physics at medium energies are listed, which are or were important for this field.

Table 7.1 Hadron accelerators for hadron physics at medium energy, LEAR and SATURNE stopped operation in 1997 and 1999, respectively.

accelerator	typ	momentum MeV/c	energy MeV	beam	cooling
TRIUMF	cyclotron	1090	500	$n, \vec{p}$	–
IUCF	synchrotron	1090	500	$A = \vec{1}, 2, 3$	e^-
SIN-PSI	cyclotron	1220	600	$\vec{p}$	–
LAMPF	linear accelerator	1460	800	$\vec{p}$	–
LEAR	synchrotron	2000	1270	$\bar{p}$	e^-
CELSIUS	synchrotron	2100	1360	$A = 1, 2, \dots$	e^-
COSY	synchrotron	3500	2620	$A = \vec{1}, 2$	e^- plus stochastic
SATURNE	synchrotron	3770	3000	$A = 1 \dots 40$	–

7.2 Targets

The choice of a certain target material for accelerator experiments as well as its chemical and mechanical properties is strongly correlated with the given experimental conditions and, certainly, with the reactions which should be investigated. Since in this article we concentrate on unpolarized meson production experiments off nucleons and deuterons, here we will restrict ourselves on corresponding targets.

Proton and deuteron targets are commonly provided using hydrogen and deuterium as raw material. In the absence of free neutron targets deuterons can be used as substitute, considering the weakly bound deuterons as effective neutron targets and restricting on reactions with the bound protons acting only as spectator particles. For the sake of completeness we want to point out that in this case the Fermi motion of the nucleons has to be considered, especially in the regime of low excess energies.

Near threshold production experiments are commonly connected with low cross sections in the order of picobarns or nanobarns. To obtain sufficient counting rates during experimental runs, high luminosities in the order of $\sim 10^{30} \text{ cm}^{-2} \text{ s}^{-1}$ are necessary and can be achieved using targets of high areal density and intense accelerator beams. Neglecting technical limitations like space charge effects, from the experimental point of view a combination of high intense ion beams and thin targets are preferable

since in this case the quality of the cooled accelerator beam is nearly uninfluenced during the passage of the target, granting well defined kinematical conditions as well as single particle interactions. In practice one has to distinguish between internal and external beam experiments. In the latter case a well prepared and extracted accelerator beam is shot onto a target and beam particles not interacting with the target are lost. Therefore, liquid or solid state targets of sufficient high areal density have commonly to be used in order to obtain the required high luminosities. Contrary, in internal beam experiments carried out at storage rings the circulating beam particles pass the target material 10^5 to 10^6 times per second and projectiles not interacting with the target remain in the beam. Therefore, to achieve similar luminosities compared to external beam experiments only thin targets are necessary. Moreover, to preserve the quality of the circulating beam and to obtain life times of the accelerated beam up to hours, effective areal target densities below $\sim 10^{15}$ atoms/cm² are desirable and can be achieved using gas and cluster jets, pellets and thin fibers. Considering hydrogen or deuterium as raw material such targets can be realized in different ways.

7.2.1 Fiber Targets

Fiber targets with diameters of only a few micrometers are of special interest for storage ring experiments with the request of high target densities in combination with a point-like target and UHV conditions. Due to the relatively simple target installation, 4π detection systems are possible. Using carbon for example as target material, fiber diameters down to $5\mu\text{m}$ are accessible, corresponding to $\sim 2 \cdot 10^{14}$ atoms/cm² [409] and allowing to compensate beam heating effects by beam cooling devices. Fiber targets for interactions off protons and deuterons naturally consist of chemical compounds like propylene such as the target for the internal beam experiment EDDA at COSY, where propylene fibers with diameters of $\sim 5\mu\text{m}$ are used. However, apart from those advantages one has to keep in mind that the mass of such CH₂ fibers is mostly given by “carrier” material carbon, which influences the performance of the data acquisition system and, moreover, causes a physical background in the data which has to be considered.

7.2.2 Liquid Targets

Liquid hydrogen and deuterium targets with their high volume densities ($\rho_{LH_2} \sim 0.07$ g/cm³) are reasonable for the use at external beam experiments. Depending on the experimental requirements large target volumes even up to several liters are accessible. However, precision experiments on the near threshold meson production force to use only small target volumes in the interaction region in order to minimize energy losses and effects of multiple scattering in the target material. An example of an experimental facility using a thin liquid hydrogen target is the COSY–TOF installation [410, 411]. Here hydrogen is liquefied using a commercial cryogenic cold head, providing a compact target volume of several millimeters in diameter. Since such targets naturally have to be encased in special vessels, the choice of appropriate entrance windows is of great importance to minimize the physical background. At COSY–TOF for example thin windows of only $\sim 1\mu\text{m}$ Mylar foil with a diameter of ~ 4 mm are routinely in use, granting good signal to background conditions.

7.2.3 Pellet Targets

Pellet target installations can be used to prepare streams of frozen gas micro-spheres as targets for internal beam experiments. Similar to fiber targets, pellet targets can be associated to the class of the solid state targets and allow to provide high areal densities in combination with point-like interaction regions, well suited for 4π detector geometries. Moreover, contrary to fiber targets these installations allow to produce pure hydrogen or deuterium pellets without any carrier material, however, at the cost of an increased experimental effort and vacuum conditions of $\sim 10^{-6}$ mbar in the scattering chamber.

At the CELSIUS ring in Uppsala such a pellet target was build up for the WASA experiment [413–415]. The pellets are produced by pressing liquid hydrogen at triple point conditions (14 K, 72 mbar) through a glass nozzle of 20 μm diameter. By means of a piezo-electric transducer connected to the nozzle the stream of hydrogen is broken into uniformly sized micro-spheres (pellets) as shown in figure 47. This device was designed to provide a stream of hydrogen pellets with diameters of 50 μm at a repetition rate of ~ 68 kHz. While the pellets itself are locally very thick ($\sim 10^{20}$ atoms/ cm^2), the effective areal density in the scattering chamber is in the order of 10^{16} atoms/ cm^2 due to the minimized geometrical beam-target overlap.




Figure 47: Formation of deuterium pellets with diameters of ~ 50 μm , the photograph was taken at the pellet target of the CELSIUS/WASA detector at CELSIUS in Uppsala.

7.2.4 Gas Jet Targets

Supersonic gas jet beams can be used as targets for both internal and external beam experiments. Produced by feeding Laval-nozzles with gases like hydrogen or deuterium, high areal target densities ($\sim 10^{16}$ atoms/ cm^2) can easily be achieved close to the nozzle exit [416]. Similar to pellet targets, proton or deuteron targets without effects of windows or carrier materials can be provided by operating the installation with gases of high purity. However, since the volume density of a gas jet beam drastically decreases both in longitudinal and transversal direction, the nozzle has to be placed very close to the interaction point in the scattering chamber. Therefore, the advantage of high target densities of such installations is commonly connected with higher gas loads in the scattering chamber and a limited spatial resolution of the interaction point if no vertex detector is used. The great advantage of such targets is the possibility to use almost all gases as target material and to adjust the areal density by

orders of magnitude by simply changing the gas input pressure.

7.2.5 Cluster Targets

Similar to conventional gas jet targets, cluster beams [417] can be produced by using fine nozzles with a convergent-divergent shape (Laval-nozzle) and, therefore, almost all gases can be used. Operated at appropriate low temperatures, a partial condensation of the used gas is possible and clusters with sizes of 10^3 to 10^6 can grow. Separated from the surrounding conventional gas beam by a set of collimators (see fig. 48), cluster beams with a homogeneous volume density, a sharp boundary, a small angular divergence and well defined beam dimensions can be prepared for interaction with circulating accelerator beams [412].




Figure 48: Schematic drawing of the principle for the formation of clusters.

Cluster beam installations are frequently used at cooler rings. Using Laval-nozzles with diameters of 11-16 μm , low nozzle temperatures of ~ 25 K and gas input pressures of ~ 18 bar, hydrogen and deuterium cluster beams with areal densities in the order of $\sim 10^{14}$ atoms/cm² in combination with UHV conditions are accessible in the interaction region. Furthermore, similar to gas beam targets the areal density can easily be adjusted over orders of magnitude by changing the gas input pressure or the nozzle temperature [412].

7.3 Typical detector arrangements

Special efforts are made for an optimal detector design to match the required conditions for kinematically completeness and undisturbed determination of the four-momentum vector of the ejectiles. A kinematically complete event is characterized by knowing all four-momentum vectors of the participating particles. The experimental information is limited by: i) statistical fluctuations, ii) the precision for the determination of the components of the four-momentum vectors, iii) contributions from any kind of background produced by means of the apparatus or due to physical reasons, iv) the efficiency of the detector covering not 100 % of the phase-space due to size or construction limitations and finally v) the reconstruction efficiency due to unavoidable inadequate performance whatsoever. Here all collaborations have to meet the optimum between wishful thinking, technical possibilities and affordable costs.

7.3.1 Example of a 4π internal-detector

The high luminosity and 4π detector CELSIUS/WASA (**W**ide **A**ngle **S**hower **A**pparatus) facility [418] is primarily




Figure 49: Top: CAD-view of the WASA detector facility

Bottom: Cross section of the central and forward part of the WASA detector facility.

A version of the article including a photo of better quality can be found at the COSY-11 homepage:

http://ikpe1101.ikp.kfa-juelich.de/cosy-11/pub/List_of_Publications.html#papers

designed for both measuring rare decay processes of light mesons and investigating meson production studies in the threshold region. The target and beam particles tend to be scattered at forward angles while the products of the meson decay will be distributed rather isotropically. The detector is constructed following these constraints and in full operation an annual production rate of about 10^{11} η 's is expected in the pp interaction.

The essential parts of the CELSIUS/WASA facility are:

- The pellet target with:

Pellets diameter and frequency of $25 - 50 \mu\text{m}$ and a few kHz, respectively, where the beam diameter is about 3 mm, 5×10^{15} atoms/cm².

- The forward detector with:

A segmented forward window counter (FWC) for fast timing in the first trigger level.

A forward proportional chamber (FPC) out of straw chambers to measure the scattering angle.

A forward trigger hodoscope (FTH) built out of three layers of circular scintillators used for:
fast trigger decisions, rough track position determination and particle identification.

A forward range hodoscope (FRH) as a plastic scintillator calorimeter for measurements of
particle energy and identification.

A forward range intermediate hodoscope (FRI) adding a position sensitivity to the FRH.

- The zero degree spectrometer to filter out low rigidity reaction products employing HPGe-
or telescopes of Si- μ -strip detectors, scintillation counters and CsI(Tl) crystals.

- The central detector with:

A scintillator electromagnetic calorimeter (SEC) consisting out of 1012 CsI(Na)
between the superconducting solenoid (SCS) and the iron yoke.

A mini drift chamber (MDC) with 1738 straw tubes for measuring tracks of charged particles.

The CELSIUS/WASA detector is completed and regular data taking has started in late 2001 on some π -production reactions where the π^0 's are detected by their $\gamma\gamma$ decays. The tagging of the η -production and the detection of the η 's by their $\gamma\gamma$ and $3\pi^0$ decays has been proven to work. In other words there is a high luminosity hadron facility ready for exciting physics.

7.3.2 Example of a centre-of-mass 4π external-detector

The COSY-TOF detection facility at the external beam line of COSY is a multi purpose modular arrangement of segmented barrel and circular planar scintillation detectors for measuring the time-of-flight and the energy loss of charged reaction products.

In Fig. 50 two CAD-drawings of a short and an extended version as well as a photograph of the outer barrel in it's longest assembly is shown. The modularity serves several purposes but essentially helps to adjust the system to the optimal dimensions according to the reaction under investigation.

Three scintillators placed 1 m, 0.5 m and 0.1 m in front of the target are used to veto a beam halo. Further the following components are employed:

- A few mm³ large liquid hydrogen/deuterium target with very thin windows [410,411]
- Different granulated start counter systems optimized for the reaction type under investigation.
- Up to three barrels equipped with granulated cylindrical detectors for timing, spatial resolution and energy loss determinations.
- Granulated three layer stop counters, divided into inner- and outer circular scintillation detector hodoscopes [419].
- A scintillation calorimeter for energy determination of charged and neutral hadrons, placed behind the stop detector, see the short version of the facility, left in fig. 50.
- In preparation a straw drift chamber hodoscope, to be installed behind the start detector systems.




Figure 50: Schematic view of the COSY-TOF detector facility in two different set-ups, the short and the longer version. Bottom: A photo from the very long version of the COSY-TOF detector.

A version of the article including a photo of better quality can be found at the COSY-11 homepage:

http://ikpe1101.ikp.kfa-juelich.de/cosy-11/pub/List_of_Publications.html#papers

7.3.3 Example of a dedicated internal threshold detector

The COSY-11 installation is an internal facility at COSY designed for threshold meson production studies. Close-to-threshold, due to the small transvers momenta, all ejectiles are limited to a narrow forward cone in the laboratory system and therefore a high geometrical acceptance is attainable with a comparatively small detector set-up. The essential components of the COSY-11 installation are:

- A cluster target located in front of a regular C-shaped COSY dipole producing H_2 or D_2 -clusters of up to 10^6 molecules and densities of about 10^{14} atoms/cm³.
- An exit window (187×7.6 cm²) of 30 μ m Al and 2×150 μ m carbon fibers to separate the ultra high beam line vacuum from atmospheric pressure. It provides high acceptance compared to conventional magnetic spectrometers especially in the horizontal plane.
- Two sets of drift chambers stacks (D1 and D2 in fig. 51) with six and eight planes, respectively and a position resolution of about 200 μ m in each plane, for particle track reconstruction and momentum analysis through the magnetic field for positively charged ejectiles.




Figure 51: Top: Top view of the COSY-11 zero degree facility.

Bottom: A photographic view of the COSY-11 detector at the COSY ring.

A version of the article including a photo of better quality can be found at the COSY-11 homepage:

http://ikp1101.ikp.kfa-juelich.de/cosy-11/pub/List_of_Publications.html#papers

- Two scintillator hodoscopes (S1 and S2 in fig. 51) for fast level trigger determination.
- A large area scintillation wall [420] ($2.2 \times 1 \text{ m}^2$) S3 in fig. 51 placed 9.1 m downstream, for fast trigger, time-of-flight and consecutively invariant mass determination.
- A detection system for negatively charged mesons consisting out of a scintillator detector and an array of silicon pad detectors, both mounted inside the dipole gap.
- A monitor system for elastically scattered particles, utilized for luminosity determination.
- In preparation for an extension: two wire chambers with a hexagonal drift structures to be positioned at the side – and at the front exit windows and a threshold Čerenkov counter behind the S1 hodoscope for pion background subtraction.

8 Outlook

Storage rings with the flexibility of different projectiles, variable beam momenta, extremely good phase-space cooling and with the unrivaled options of internal and external detection systems provide an excellent tool for investigations on meson production at threshold. Especially the high quality of relative and absolute energy resolutions available at modern accelerators are crucial for such studies.

In the present report, discussing the close-to-threshold meson production in hadronic interactions, the natural restriction to investigations using only hadronic probes was made though interesting and informative results were certainly obtained with electromagnetic tools as well.

The theory of strong interactions (QCD) has a convincing predictive power at high energies. Perturbative expansions in the coupling constant no longer converge for medium energy scales and the structure of QCD becomes increasingly complicated. The threshold meson production in nucleon–nucleon scattering offers valuable aspects for studying production, decay and interaction of hadrons and investigating the reaction mechanisms, the dynamics of the system and appropriately chosen subsystems as well as the structure of hadrons themselves. Such experiments probe the sensitivity of nucleon–nucleon interactions at distances of ≤ 1 fm, while the long-range part of the interaction is suppressed. Meson production in $pp \rightarrow ppX$ reactions (with $X = \pi^0, \eta, \omega, \phi$) is a process of strong inelasticity in proton–proton scattering.

8.1 Polarized beams and/or targets

In the present paper discussions utilizing polarized beam and/or targets are omitted. At the upgraded IUCF cooler ring a polarized proton beam in combination with a windowless polarized hydrogen target now permits measurements of all spin correlation coefficients for $\vec{p}\vec{p} \rightarrow NN\pi$ reactions. Very recently initial results for spin correlations were reported [421]. The goal of such experiments is to qualify the energy dependence and the importance of higher partial waves by determining analyzing powers and spin correlations. Here the hadron physics community certainly faces the challenge of new experimental techniques with necessary improvements of theoretical developments, since polarization observables are sensitive indicators of reaction mechanisms, relative contribution of partial waves and transition amplitudes.

The production of η mesons near threshold has been measured by different groups [61–63, 65]. Very close-to-threshold only S-waves are supposed to contribute to the production process and in fact the S-wave production is large due to the presence of the $N^*(1535)S_{11}$ resonance which seems to act as an intermediate state.

Both TOF [98] and COSY-11 [66] took recently rather high statistics data on the η production at $Q \approx 15.5$ MeV with fairly flat angular distributions as expected for an s-wave production. Still – whether some data suggest or don't suggest – differential cross section measurements might be too insensitive to reveal e.g. a small Sd -contribution. Data using a polarized beam, however, might indicate effects arising from an interference term of the kind:

$$2\text{Re}\{A_{Ss}A_{Sd}^*\} \cos 2\theta_\eta^* \quad (56)$$

where A_{Ss} is the amplitude for the production of a final pp pair with angular momentum $L_{pp} = 0$ and $l = 0$ while for A_{Sd} the η meson has an angular momentum of $l_\eta = 2$.

The importance of d-waves in η production is well known from both $\pi^-p \rightarrow \eta n$ [422] and $\gamma p \rightarrow \eta p$ [397] reactions and the fact that this interference term is negative suggests that the η -production in proton–proton scattering is governed mainly by ρ -exchange [99]. It is the aim of the COSY-11 group, followed by a recent feasibility study [181] to investigate the presence of this and other higher partial waves through the study of the analyzing power A_y .

Though the asymmetry should also be studied with respect to the final proton directions, the s - d interference contributes significantly to the η analyzing power defined as:

$$A_y := \frac{\text{Tr}(\mathcal{M} \sigma_y \mathcal{M}^\dagger)}{\text{Tr}(\mathcal{M} \mathcal{M}^\dagger)} \quad (57)$$

with the transition matrix element $\mathcal{M}$ and the Pauli matrix σ_y .

To lowest order in the η d -wave amplitude, the analyzing power is proportional to the imaginary part of the s - d interference [423],

$$A_y \approx 2 \frac{\text{Im}\{A_{Ss} A_{Sd}^*\}}{|A_{Ss}|^2} \quad (58)$$

where $\phi_\eta^* = 0$ corresponds to the plane of the beam (perpendicular to the polarization direction). On angular momentum grounds, this signal is expected to vary like $(p_\eta^{cm})^2$, *i.e.* roughly like Q .

The $P_{11}(1440)$ resonance could also be of importance for the η production [424], leading to p -waves in the final state. The interference between the Ps and Pp amplitudes gives an analyzing power proportional to:

$$A_y \approx 2 \frac{\text{Im}\{A_{Ps} A_{Pp}^*\}}{|A_{Ss}|^2} \quad (59)$$

and such a signal has been observed in the $pp \rightarrow pp\pi^0$ reaction at low energies [38]. The presence of the three p -waves in the interference might suppress such a term close-to-threshold for η production and no unambiguous signal for them has been seen in the unpolarized differential cross section. Nevertheless one must be prepared for their existence. The Ss - Sd interference gives a maximal A_y signal for $\theta_\eta^* \approx 45^\circ$ and vanishes at 90° , whereas the Ps - Pp signal should be largest at this point.

Both the differential cross section and the analyzing power should be studied in order to detect effects associated with the onset of higher partial waves. It should be noted that the effects of incomplete acceptance are much less of a problem for A_y than for the $d\sigma/d\Omega$ measurement provided that the beam characteristics of the two polarization states are similar. Finally, the results should help to settle the on-going discussion of whether the η production is dominated by ρ [175, 176, 425, 426], ω [178] or η [19] exchange.

First data on the η meson production with a polarized proton beam in the reaction have been taken at the internal experiment facility COSY-11 [181]. The measurement was performed with a proton beam momentum $\vec{P}_p = 2.096 \text{ GeV}/c$ corresponding to an excess energy of $Q = 40 \text{ MeV}$. In Figure 52 a comparison of the data (solid triangles) with theoretical predictions for $Q = 37 \text{ MeV}$ from Fäldt and Wilkin [177] (dotted line) and Nakayama [20] (solid line) is shown. In addition, the theoretical calculations for $Q = 10 \text{ MeV}$ are plotted, both based on the one meson exchange model. While Fäldt and Wilkin predict a dominance of ρ exchange (dashed line), Nakayama concludes a dominant π and η exchange (solid line).




Figure 52: Theoretical predictions for the analyzing power according to [177] (dotted line) and [20] (solid line) for $Q = 10$ and 37 MeV. The solid triangles represent the measured values [181] at $Q = 40$ MeV.

Though the data are consistent with $A_y = 0$ within the error bars given, there seems to be a slight increase of A_y towards backward scattering angles. First results of the analyzing power – deduced from the very first feasibility study run – suffer from statistics but tend to differ from both theoretical calculations, although the predictions of [177] show a quite good match. For a more precise statement further measurements are needed which could benefit from higher polarization now available at COSY.

8.2 Final state interaction

An important physics aspect of the modern cooler ring facilities is the interaction of particles in the final state. A detailed study of the Dalitz plot of, e.g. the reaction $pp \rightarrow pK^+\Lambda$ measured at COSY-11 [328], made possible the extraction of valuable information about the Λ - N interaction.

The $pn \rightarrow d\eta$ reaction, measured at TSL in Uppsala [47–50], limited significantly the range of values for the η - N scattering length. This quantity is the relevant input for calculations studying the existence of η -nucleus bound states.

8.3 Baryon resonances

Throughout the article the importance of an intermediate excitation of baryon resonances has been pointed out.

At excess energies of 1–2 GeV, baryon resonances begin to play an important role in nucleon–nucleon collisions. Most of our understanding about these resonances comes from pion and photon induced reactions on the nucleon. Current quark models predict many more resonances than can be experimentally extracted from the data. A large program at both JLAB and ELSA focuses on this so-called missing resonance problem and the search for yet unknown baryonic states in electro-nuclear reactions. However, besides the possibility that resonances thus far simply went unnoticed because the quality of the data was insufficient, another explanation might be that these resonances couple only weakly to photons or pions. An accelerator like COSY, built for nucleon–nucleon as well as nucleon–nucleus collisions will provide for the necessary experiments to fill that gap in the future.

8.4 Symmetries

The investigation of symmetries and their breaking has always led to significant insights into the inner workings of nature. Already a large program at COSY [405, 406] is studying charge-symmetry breaking

- an effect closely linked to the difference in the current masses of the light quarks.

Another interesting aspect of charge-symmetry breaking is its high sensitivity: Once, for instance, the f_0 - a_0 mixing amplitude is isolated, it will provide us with insights about the nature of scalar resonances. Studies about the structure of these resonances are under investigations at COSY [278–280].

At the CELSIUS ring in Uppsala the CELSIUS/WASA detection system is in operation now and exciting experiments especially on η -meson decay studies will start.

8.5 Meson production in the three-nucleon system

Naively, one would think that near-threshold meson production in pd collisions can be fully understood in terms of a short-ranged $NN \rightarrow NN\pi$ process, with the extra nucleon acting as a spectator. However, backward π production in the $pd \rightarrow {}^3He\pi^0$ cannot be explained in terms of a spectator model. In the case of meson production in pd collisions the number of participating amplitudes is small as well, even though the transferred momentum is large. Hereby a unique environment is available for studying the interaction of more than just two nucleons, e.g. three-nucleon forces. It is likely that eventually the research on three-nucleon forces will focus on this process. Again, the role of polarization observables is important.

8.6 Apparative aspects

For the experiments, the development of suitable experimental configurations, comprising polarized targets and associated detector systems that facilitate measurements of polarization observables is necessary.

Present and future experiments become possible only through the outstanding performance of the cooler accelerators. With the new injector for COSY the situation will be further improved. It will be possible to fill the machine up to the space-charge limit with polarized protons and deuterons, whereby electron-cooled beams at injection energy and stochastically-cooled beams at high energy of unprecedented brilliance will be available to the users. Efficient correction schemes to preserve the polarization of protons during acceleration have already been established, while for deuterons no difficulties are anticipated, because no depolarizing resonances occur in the energy range of COSY. New polarization experiments in the future will certainly call for further refinement of the spin-manipulation systems in COSY, e.g. longitudinally polarized internal and external beams, for which suitable spin rotators have to be developed.

8.7 Epilogue

The above mentioned examples – which will be the subject of a Summer School and Workshop on COSY Physics [427] – demonstrate the richness of physics in meson production reactions. The drawback of this wealth comes at the expense that, apart from rare cases, it is difficult to extract a particular piece of information from the data, e.g. both resonances and final state interactions modify the invariant mass plot in a coherent superposition.

Fortunately, the use of polarization in the initial state acts as a spin filter and different contributions to the interaction can be singled out, because they show up in the angular distributions of different spin combinations. It is important to stress, that as long as only the lowest partial waves contribute significantly to the process under study, the number of independent observables from experiments with polarized beam and target is sufficient to allow for a model independent partial wave analysis.

Acknowledgements

The authors would like to express their utmost gratitude to the numerous people who gave the various help needed in order to successfully complete this article.

Especially we would like to thank R. Czyżykiewicz, J. Flood, J. Haidenbauer, Ch. Hanhart, T. Sefzick and P. Winter for both scientific and technical support throughout the preparation of this manuscript.

References

- [1] V. Bernard, N. Kaiser and U.-G. Meißner, *Int. J. Mod. Phys. E* 4 (1995) 193
- [2] Proceedings of the *Workshop On Chiral Dynamics: Theory And Experiment*, July 25–29, 1994, Cambridge, Massachusetts; Eds. A.M. Bernstein and B.R. Holstein (Springer, Berlin), *Lecture Notes in Physics* 452 (1995)
- [3] C. Wilkin, *e-Print Archive*: nucl-th/9810047 and Proceedings of the *8th International Conference On The Structure Of Baryons (Baryons 98)*, September 22–26, 1998, Bonn, Germany; Eds. D.W. Menze and B. Metsch (World Scientific, Singapore, 1999), p.505
- [4] M. Gell-Mann and K.M. Watson, *Ann. Rev. Nucl. Sci.* 4 (1954) 219
- [5] A.H. Rosenfeld, *Phys. Rev.* 96 (1954) 130
- [6] D. Koltun and A. Reitan, *Phys. Rev.* 141 (1966) 1413
- [7] M.E. Schillaci et al., *Phys. Rev.* 179 (1969) 1539
- [8] H.O. Meyer et al., *Nucl. Phys. A* 539 (1992) 633
- [9] H.O. Meyer et al., *Phys. Rev. Lett.* 65 (1990) 2846
- [10] W.W. Daehnick et al., *Phys. Rev. Lett.* 74 (1995) 2913
- [11] T.-S.H. Lee and D.O. Riska, *Phys. Rev. Lett.* 70 (1993) 2237
- [12] C. Hanhart et al., *Phys. Lett. B* 358 (1995) 21
- [13] C. Hanhart et al., *Phys. Lett. B* 444 (1998) 25
- [14] C. Hanhart et al., *Phys. Rev. C* 61 (2000) 064008
- [15] B. Blankleider and A.N. Kvinikhidze, *Few Body Systems Suppl.* 12 (2000) 223
- [16] S. Wycech and A.M. Green, *Nucl. Phys. A* 663 (2000) 529
- [17] A. Fix and H. Arenhövel, *Nucl. Phys. A* 697 (2002) 277
- [18] K.M. Watson, *Phys. Rev.* 88 (1952) 1163
- [19] M. Batinić, A. Švarc, T.-S.H. Lee, *Phys. Scripta* 56 (1997) 321
- [20] K. Nakayama, *e-Print Archive* nucl-th/0108032 and Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 119
- [21] C. Hanhart and K. Nakayama, *Phys. Lett. B* 454 (1999) 176
- [22] F. Kleefeld, *e-Print Archive* nucl-th/0108064 and Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 51
- [23] C. Hanhart, U. van Kolck and G.A. Miller, *Phys. Rev. Lett.* 85 (2000) 2905 and references therein
- [24] N.I. Kochelev, V. Vento and A.V. Vinnikov, *Phys. Lett. B* 472 (2000) 247

- [25] M. Dillig, *e-Print Archive* hep-ph/0202067
- [26] N. Kaiser, P.B. Siegel and W. Weise, *Nucl. Phys. A* 594 (1995) 325
- [27] N. Kaiser, T. Waas and W. Weise, *Nucl. Phys. A* 612 (1997) 297
- [28] W. Weise and U.-G. Meißner, HESR-Project, Contributions to the Scientific Case, (1999) 103
- [29] D. Harrach, Proceedings of the *29th International Workshop On Gross Properties Of Nuclei and Nuclear Excitations (Hirschegg 2001: Structure Of Hadrons)*, January 14–20, 2001, Hirschegg, Austria; Eds. H. Feldmeier, J. Knoll, W. Nörenberg and J. Wambach (GSI, Darmstadt, 2001)
- [30] R.L. Workman, in *Eur. Phys. J. C* 15 (2000) 698
- [31] R. Machleidt and I. Slaus, *J. Phys. G* 27 (2001) R69
- [32] R. Machleidt, *Phys. Rev. C* 63 (2001) 024001
- [33] A. Moalem, E. Gedalin, L. Razdolskaja, and Z. Shorer, *Nucl. Phys. A* 600 (1996) 445
- [34] V. Bernard, N. Kaiser, U.-G. Meißner, *Eur. Phys. J. A* 4 (1999) 259
- [35] E. Gedalin, A. Moalem, L. Razdolskaja, *Nucl. Phys. A* 650 (1999) 471
- [36] E. Byckling and K. Kajantie, *Particle Kinematics*, (John Wiley & Sons, New York, 1973)
- [37] P. Moskal et al., *Phys. Lett. B* 482 (2000) 356
- [38] H.O. Meyer et al., *Phys. Rev. Lett.* 81 (1998) 3096
- [39] D. Grzonka and K. Kilian, Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 100
- [40] R.H. Dalitz, *Phil. Mag.* 44 (1953) 1068
- [41] J. Dyring, *Ph.D. thesis*, University of Uppsala, *Acta Universitatis Upsaliensis* 14 (1997)
- [42] D. Chodrow, *Nuovo Cimento* 50 (1967) 674
- [43] D.A. Hutcheon et al., *Phys. Rev. Lett.* 64 (1990) 176
- [44] D.A. Hutcheon et al., *Nucl. Phys. A* 535 (1991) 618
- [45] M.G. Bachman et al., *Phys. Rev. C* 52 (1995) 495
- [46] R. Helmer, *Can. J. Phys.* 65 (1987) 588
- [47] R. Bilger et al., *Nucl. Instr. and Meth. A* 457 (2001) 64
- [48] H. Calén et al., *Phys. Rev. Lett.* 80 (1998) 2069
- [49] H. Calén et al., *Phys. Rev. C* 58 (1998) 2667
- [50] H. Calén et al., *Phys. Rev. Lett.* 79 (1997) 2642
- [51] P. Moskal, *e-Print Archive*: nucl-ex/0110001 and Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 27
- [52] R.J. Slobodrian, *Rep. Prog. Phys.* 34 (1971) 175
- [53] M. Lacombe et al., *Phys. Lett. B* 101 (1981) 139
- [54] S. Haggström, *Ph.D. thesis*, Uppsala University, *Acta Universitatis Upsaliensis* 13 (1997)
- [55] M. Lacombe et al., *Phys. Rev. C* 21 (1980) 861
- [56] M. Büscher et al., COSY Proposals No. 75 (1998) and No. 94 (2000), <http://www.fz-juelich.de/ikp/anke>
- [57] F. Duncan et al., *Phys. Rev. Lett.* 80 (1998) 4390
- [58] M. Garçon, J.W. Van Orden, *e-Print Archive*: nucl-th/0102049, submitted to *Adv. Nucl. Phys.*

- [59] E. Chiavassa et al., *Phys. Lett. B* 337 (1994) 192
- [60] R. Smith and C. Wilkin, *Nuovo Cimento Lett. IV* (1970) 647
- [61] J. Smyrski et al., *Phys. Lett. B* 474 (2000) 182
- [62] F. Hibou et al., *Phys. Lett. B* 438 (1998) 41
- [63] H. Calén et al., *Phys. Lett. B* 366 (1996) 39
- [64] E. Chiavassa et al., *Phys. Lett. B* 322 (1994) 270
- [65] A.M. Bergdolt et al., *Phys. Rev. D* 48 (1993) R2969
- [66] P. Moskal et al., *e-Print Archive: nucl-ex/0110018* and *πN Newsletter* 16 (2002) 367
- [67] J. Bystricky and F. Lehar, *Physics Data* Nr.11-1, Fachinfo.zentrum, Karlsruhe, 1987, Pt. I, p.98
- [68] H. Rohdjesse et al., *Phys. Rev. Lett.* 70 (1993) 2864
- [69] J. Greiff et al., *Phys. Rev. C* 62 (2000) 064002
- [70] H.O. Meyer and J.A. Niskanen, *Phys. Rev. C* 47 (1993) 2474
- [71] P. Moskal et al., *Phys. Lett. B* 517 (2001) 295
- [72] P. Moskal et al., *e-Print Archive: nucl-ex/0201016* and *J. Phys. G* 28 (2002) 1777
- [73] P. Senger et al., *Prog. Part. Nucl. Phys.* 42 (1999) 209
- [74] J. Schaffner-Bielich, I.N. Mishustin and J. Bondorf, *Nucl. Phys. A* 625 (1997) 325
- [75] S. Goldhaber et al., *Phys. Rev. Lett.* 9 (1962) 135 ,
W. Cameron et al., *Nucl. Phys. B* 78 (1974) 93 , C.J. Adams et al., *Nucl. Phys. B* 66 (1973) 36
- [76] C.J. Adams et al., *Nucl. Phys. B* 96 (1975) 54 , R. Armenteros et al., *Nucl. Phys. B* 21 (1970) 15 ,
T.S. Mast et al., *Phys. Rev. D* 14 (1976) 13 , M. Sakitt et al., *Phys. Rev.* 139 (1965) B719
- [77] C. Quentmeier et al., *Phys. Lett. B* 515 (2001) 276
- [78] F. Balestra et al., *Phys. Lett. B* 468 (1999) 7
- [79] A. Sibirtsev, W. Cassing and C.M. Ko, *Z. Phys. A* 358 (1997) 101
- [80] J. Złomańczuk et al., *Phys. Lett. B* 436 (1998) 251
- [81] R. Bilger et al., *Nucl. Phys. A* 693 (2001) 633
- [82] H.O. Meyer et al., *Phys. Rev. C* 63 (2001) 064002
- [83] J.D. Jackson, J.M. Blatt, *Rev. Mod. Phys.* 22 (1950) 77
- [84] H.A. Bethe, *Phys. Rev.* 76 (1949) 38
- [85] D.Y. Wong and H.P. Noyes, *Phys. Rev.* 126 (1962) 1866
- [86] R.A. Arndt, R.H. Hackman and L.D. Roper, *Phys. Rev. C* 15 (1977) 1002
- [87] E. Oset, A. Ramos, *Nucl. Phys. A* 635 (1998) 99
- [88] J. Caro Ramon, N. Kaiser, S. Wetzel, W. Weise, *Nucl. Phys. A* 672 (2000) 249
- [89] A. Bondar et al., *Phys. Lett. B* 356 (1995) 8
- [90] P. Moskal et al., *Phys. Lett. B* 474 (2000) 416
- [91] M. Wolke, C. Quentmeier et al., COSY Proposal No. 61.2 (2000), P. Winter, M. Siemaszko et al.,
COSY Proposal No. 61.3 (2001), <http://ikpe1101.ikp.kfa-juelich.de>
- [92] M. Wolke and A. Sibirtsev, Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke,
Schriften des FZ-Jülich – Matter and Materials 11 (2002) 239
- [93] R.A. Arndt et al., *Phys. Rev. C* 56 (1997) 3005
The Virginia Tech Partial-Wave Analysis Facility (SAID) <http://gwdac.phys.gwu.edu>
- [94] H.O. Meyer et al., *Phys. Rev. Lett.* 83 (1999) 5439

- [95] J.R. Bergervoet et al., *Phys. Rev. C* 41 (1990) 1435
- [96] L.D. Knutson, in *Nuclear Physics at Storage Rings*, edited by H.O. Meyer and P. Schwandt, *AIP Conf. Proc.* 512 (2000) 177
- [97] F. Balestra et al., *Phys. Lett. B* 491 (2000) 29
- [98] M. Abdel-Bary et al., e-Print Archive: nucl-ex/0205016, to be published in *Eur. Phys. J.*
- [99] H. Calén et al., *Phys. Lett. B* 458 (1999) 190
- [100] A.B. Migdal, *Sov. Phys. JETP* 1 (1955) 2
- [101] B.J. Morton et al., *Phys. Rev.* 169 (1968) 825
- [102] J.P. Naisse, *Nucl. Phys. A* 278 (1977) 506
- [103] H.P. Noyes and H.M. Lipinski, *Phys. Rev. C* 4 (1971) 995
- [104] H.P. Noyes, *Ann. Rev. Nucl. Sci.* 22 (1972) 465
- [105] V.G.J. Stoks et al., *Phys. Rev. C* 48 (1993) 792
- [106] B.L. Druzhinin, A.E. Kudryavtsev and V.E. Tarasov, *Z. Phys. A* 359 (1997) 205
- [107] R. Shyam and U. Mosel, *Phys. Lett. B* 426 (1998) 1
- [108] M.L. Goldberger and K.M. Watson, *Collision Theory*, (John Wiley & Sons, New York, 1964)
- [109] S. Stanislaus et al., *Phys. Rev. C* 41 (1990) R1913
- [110] G. Fäldt and C. Wilkin, *Phys. Lett. B* 382 (1996) 209
- [111] P. Moskal et al., *Phys. Rev. Lett.* 80 (1998) 3202
- [112] A. Khoukaz et al., *Annual Report 2001*, Institut für Kernphysik, Universität Münster, in press
- [113] J.T. Balewski et al., *Phys. Lett. B* 420 (1998) 211
- [114] S. Sewerin et al., *Phys. Rev. Lett.* 83 (1999) 682
- [115] P. Kowina, *Ph.D. thesis*, Silesian University Katowice (2002), in preparation
- [116] R. Bilger et al., *Phys. Lett. B* 420 (1998) 217
- [117] S. Marcello et al., *Nucl. Phys. A* 691 (2001) 344c
- [118] D. Hesselbarth, *Ph.D. thesis*, University of Bonn, *Berichte des FZ-Jülich*, Jül-3849 (2001)
- [119] W.J. Fickinger et al., *Phys. Rev.* 125 (1962) 2082
- [120] G. Fäldt and C. Wilkin, *Phys. Rev. C* 56 (1997) 2067
- [121] A. Moalem et al., *Nucl. Phys. A* 589 (1995) 649 , e-Print Archive: hep-ph/9505264
- [122] A.M. Green and S. Wycech, e-Print Archive: nucl-th/0009053
- [123] A. Sibirtsev et al., *Phys. Rev. C* 65 (2002) 044007
- [124] D. Sigg et al., *Nucl. Phys. A* 609 (1996) 269 , *Nucl. Phys. A* 617 (1997) 526
- [125] A.M. Green and S. Wycech, *Phys. Rev. C* 55 (1997) R2167
- [126] W. Glöckle et al., *Phys. Rep.* 274 (1996) 107
- [127] U. Schuberth, *Ph.D. thesis*, University of Uppsala, *Acta Universitatis Upsaliensis* 5 (1995)
- [128] V. Baru et al., *Phys. Atom. Nucl.* 64 (2001) 579
- [129] J.A. Niskanen, *Phys. Lett. B* 456 (1999) 107
- [130] H.W. Fearing, *Phys. Rev. Lett.* 81 (1998) 758
- [131] J. Haidenbauer et al., *Phys. Rev. C* 48 (1993) 2190
- [132] H. Van Haeringen, *Nucl. Phys. A* 253 (1975) 355
- [133] K. Nakayama et al., *Phys. Rev. C* 61 (2000) 024001
- [134] D.E. Groom et al., *Eur. Phys. J. C* 15 (2000) 1

- [135] M. Batinić et al., *e-Print Archive*: nucl-th/9703023
- [136] J. Smyrski, Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 46
- [137] A. Fix and H. Arenhövel, *Eur. Phys. J. A* 9 (2000) 119
- [138] S.A. Rakityansky et al., *Phys. Rev. C* 53 (1996) R2043
- [139] A. Švarc and S. Ceci, Contribution to N*2000, *e-Print Archive*: nucl-th/0009024
- [140] C. Wilkin, *Phys. Rev. C* 47 (1993) R938
- [141] N.V. Shevchenko et al., *Eur. Phys. J. A* 9 (2000) 143
- [142] S. Wycech, *Acta Phys. Pol. B* 27 (1996) 2981
- [143] H. Garcilazo and M.T. Peña, *Phys. Rev. C* 63 (2001) 021001
- [144] M. Green, S. Wycech, *Phys. Rev. C* 60 (1999) 035208
- [145] T. Ueda, *Phys. Rev. Lett.* 66 (1991) 297, *Phys. Lett. B* 291 (1992) 228
- [146] T. Ueda, *Phys. Scripta* 48 (1993) 68
- [147] S. Wycech and A.M. Green, *Phys. Rev. C* 64 (2001) 045206
- [148] A. Deloff, *Phys. Rev. C* 61 (2000) 024004
- [149] F. Hibou et al., *Eur. Phys. J. A* 7 (2000) 537
- [150] A. Gillitzer et al., COSY Proposal No. 102 (2001) <http://www.fz-juelich.de/ikp/COSY-TOF>
- [151] G. Rappenecker et al., *Nucl. Phys. A* 590 (1995) 763
- [152] W.B. Tippens et al., *Phys. Rev. D* 63 (2001) 052001
- [153] P. Moskal et al., *Annual Report 2000*, IKP, FZ-Jülich, Jül-3852 (2001) 48
- [154] P. Moskal, *Ph.D. thesis*, Jagellonian University, Cracow, *Berichte des FZ-Jülich*, Jül-3685 (1999) <http://ikpe1101.ikp.kfa-juelich.de>
- [155] S. Capstick et al., *e-Print Archive*: hep-ph/0012238 and *Hadronic Physics White Paper*, proceedings of the *APS Division of Nuclear Physics Town Meeting on Electromagnetic and Hadronic Physics*, December 1–4, 2000, Newport News, Virginia; <http://www.jlab.org/intralab/calendar/dnp-tm>
- [156] K. Maltman and N. Isgur, *Phys. Rev. D* 29 (1984) 952
- [157] J.A. Niskanen, *Phys. Lett. B* 289 (1992) 227
- [158] J. Haidenbauer, C. Hanhart and J. Speth, *Acta Phys. Pol. B* 27 (1996) 2893
- [159] J.A. Niskanen, *Phys. Rev. C* 49 (1994) 1285
- [160] E. Hernández and E. Oset, *Phys. Lett. B* 350 (1995) 158
- [161] C.J. Horowitz, H.O. Meyer and D.K. Griegel, *Phys. Rev. C* 49 (1994) 1337
- [162] H. Machner and J. Haidenbauer, *J. Phys. G* 25 (1999) R231
- [163] J. Adam et al., *Phys. Lett. B* 407 (1997) 97
- [164] T. Sato et al., *Phys. Rev. C* 56 (1997) 1246
- [165] T.D. Cohen et al., *Phys. Rev. C* 53 (1996) 2661
- [166] B.-Y. Park et al., *Phys. Rev. C* 53 (1996) 1519
- [167] U. van Kolck, G.A. Miller, D.O. Riska, *Phys. Lett. B* 388 (1996) 679
- [168] S.K. Saha et al., *Phys. Lett. B* 461 (1999) 175
- [169] B. v. Przewoski et al., *Phys. Rev. C* 61 (2000) 064604

- [170] T. Feldmann and P.Kroll, *e-Print Archive*: hep-ph/0201044 and
Proceedings of the *Workshop on Eta Physics: Prospects Of Precision Measurements With The CELSIUS/WASA Facility*, October 25–27, 2001, Uppsala, Sweden; to be published in *Phys. Scripta*
- [171] W. Grein, P. Kroll, *Nucl. Phys. A* 338 (1980) 332
- [172] M.M. Nagels et al., *Nucl. Phys. B* 147 (1979) 189 , *Few body syst. and nucl. forces I* (1978) 17
- [173] K. Nakayama, J. Speth and T.-S.H. Lee, *Phys. Rev. C* 65 (2002) 045210
- [174] R. Machleidt, *Adv. Nucl. Phys.* 19 (1989) 189
- [175] J.F. Germond and C. Wilkin, *Nucl. Phys. A* 518 (1990) 308
- [176] J.M. Laget, F. Wellers, J.F. Lecomte, *Phys. Lett. B* 257 (1991) 254
- [177] G. Fäldt and C. Wilkin, *Phys. Scripta* 64 (2001) 427
- [178] T. Vetter, A. Engel, T. Biró and U. Mosel, *Phys. Lett. B* 263 (1991) 153
- [179] B.L. Alvaredo and E. Oset, *Phys. Lett. B* 324 (1994) 125
- [180] M.T. Peña, H. Garcilazo and D.O. Riska, *Nucl. Phys. A* 683 (2001) 322
- [181] P. Winter, *diploma thesis*, University of Bonn,
Berichte des FZ-Jülich, Jül-3943 (2002), <http://ikpe1101.ikp.kfa-juelich.de>
- [182] A. Sibirtsev and W. Cassing, *Eur. Phys. J. A* 2 (1998) 333
- [183] R. Plötzke et al., *Phys. Lett. B* 444 (1998) 555
- [184] P.Ball, J.-M. Frère and M. Tytgat, *Phys. Lett. B* 365 (1996) 367
- [185] S.D. Bass, *Phys. Lett. B* 463 (1999) 286
- [186] S.D. Bass, S. Wetzels and W. Weise, *Nucl. Phys. A* 686 (2001) 429
- [187] S.D. Bass, *e-Print Archive*: hep-ph/0006348 and Proceedings of the *8th Int. Workshop On Deep Inelastic Scattering And QCD (DIS 2000)*, April 25–30, 2000, Liverpool, England;
Eds. J.A. Gracey and T. Greenshaw (World Scientific, Singapore, 2001), p.206
- [188] S.D. Bass, *Eur. Phys. J. A* 5 (1999) 17
- [189] N. Nikolaev, *CosyNews* 3 (1998) Published by the FZ-Jülich in Cooperation with CANU.
- [190] E. Kou, *Phys. Rev. D* 63 (2001) 054027
- [191] S.D. Bass, *private communication*
- [192] S. D. Bass, *e-Print Archive*: hep-ph/0108187 and
Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*,
June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke,
Schriften des FZ-Jülich – Matter and Materials 11 (2002) 11
- [193] F. Kleefeld, M. Dillig and F. Pilotto, *Acta Phys. Pol. B* 27 (1996) 2867
- [194] M. Dillig, *e-Print Archive*: nucl-th/0201050
- [195] S. Okubo, *Phys. Lett.* 5 (1963) 165 , G. Zweig, *Report CERN-TH-412* (1964)
J. Iizuka, *Prog. Theor. Phys. Suppl.* 37–38 (1966) 21
- [196] H. Lipkin, *Int. J. Mod. Phys. E* 1 (1992) 603
- [197] H. Lipkin, *Phys. Lett. B* 60 (1976) 371
- [198] T.P. Cheng and R.F. Dashen, *Phys. Rev. Lett.* 26 (1971) 594 , T.P. Cheng, *Phys. Rev. D* 13 (1976) 2161 , J.F. Donoghue and C.R. Nappi, *Phys. Lett. B* 168 (1986) 105 , J. Gasser, H. Leutwyler, M.P. Locher and M.E. Sainio, *Phys. Lett. B* 213 (1988) 85 , J. Gasser, H. Leutwyler and M.E. Sainio, *Phys. Lett. B* 253 (1991) 252
- [199] J. Ashman et al., *Phys. Lett. B* 206 (1988) 364 , J. Ashman et al., *Nucl. Phys. B* 328 (1989) 1 ,
P. Amaudruz et al., *Phys. Lett. B* 295 (1992) 159

- [200] T. Adams et al., *e-Print Archive*: hep-ex/9906038 and
 Proceedings of the *Workshop on Physics with Electron Polarized Ion Collider (EPIC '99)*, April
 8–11, 1999, Bloomington, Indiana; Eds. L.C. Bland, J.T. Londergan and A.P. Szczepaniak (World
 Scientific, Singapore, 2000), p.337
- [201] V. Barone, C. Pascaud and F. Zomer, *Eur. Phys. J. C* 12 (2000) 243 and references therein
- [202] J. Ellis, M. Karliner, D.E. Kharzeev and M.G. Sapozhnikov, *Phys. Lett. B* 353 (1995) 319
- [203] T. Gutsche, A. Faessler, G.D. Yen and S.N. Yang, *Nucl. Phys. B Proc. Suppl.* 56 (1997) 311
- [204] C.B. Dover, T. Gutsche, M. Maruyama and A. Faessler, *Prog. Part. Nucl. Phys.* 29 (1992) 87
- [205] L. Bertolotto et al., *Phys. Lett. B* 345 (1995) 325 ,
 C. Evangelista et al., *Phys. Rev. D* 57 (1998) 5370
- [206] M.P. Locher, Y. Lu and B.S. Zou, *Z. Phys. A* 347 (1994) 281
- [207] M.P. Locher and Y. Lu, *Z. Phys. A* 351 (1995) 83
- [208] V. Mull, K. Holinde and J. Speth, *Phys. Lett. B* 334 (1994) 295
- [209] D. Buzatu and F.M. Lev, *Phys. Lett. B* 329 (1994) 143 , *Phys. Rev. C* 51 (1995) 2893 ,
Phys. Part. Nucl. 29 (1998) 88
- [210] O. Gortchakov, M.P. Locher, V.E. Markushin and S. von Rotz, *Z. Phys. A* 353 (1996) 447
- [211] A.V. Anisovich and E. Klempt, *Z. Phys. A* 354 (1996) 197
- [212] U.-G. Meißner, V. Mull, J. Speth and J.W. van Orden, *Phys. Lett. B* 408 (1997) 381
- [213] V.E. Markushin and M.P. Locher, *Eur. Phys. J. A* 1 (1998) 91
- [214] P. Geiger and N. Isgur, *Phys. Rev. D* 55 (1997) 299
- [215] A.M. Bertin et al., *Phys. Lett. B* 388 (1996) 450
- [216] F. Balestra et al., *Phys. Rev. Lett.* 81 (1998) 4572 , *Phys. Rev. C* 63 (2001) 024004
- [217] V. Blobel et al., *Phys. Lett. B* 59 (1975) 88 , R. Baldi et al., *Phys. Lett. B* 68 (1977) 381 ,
 M.W. Arenton et al., *Phys. Rev. D* 25 (1982) 2241 ,
 S.V. Golovkin et al., *Z. Phys. A* 359 (1997) 435
- [218] A. Sibirtsev, *Nucl. Phys. A* 604 (1996) 455
- [219] A. Sibirtsev and W. Cassing, *Eur. Phys. J. A* 7 (2000) 407
- [220] W.S. Chung, G.Q. Li and C.M. Ko, *Phys. Lett. B* 401 (1997) 1
- [221] A.I. Titov, B. Kämpfer and V.V. Shklyar, *Phys. Rev. C* 59 (1999) 999
- [222] A. Sibirtsev and W. Cassing, *Few Body Systems Suppl.* 99 (2000) 1
- [223] S. Abd El-Samad et al., *Phys. Lett. B* 522 (2001) 16
- [224] M.P. Rekalo, J. Arvieux and E. Tomasi-Gustafsson, *Z. Phys. A* 357 (1997) 133
- [225] K. Nakayama et al., *Phys. Rev. C* 57 (1998) 1580
- [226] K. Nakayama et al., *Phys. Rev. C* 60 (1999) 055209
- [227] G. Höhler et al., *Nucl. Phys. B* 114 (1976) 505
- [228] S. Furuichi and K. Watanabe, *Prog. Theor. Phys.* 82 (1989) 581 and 83 (1990) 565
- [229] R. Machleidt, K. Holinde and C. Elster, *Phys. Rep.* 149 (1987) 1
- [230] P. Mergell, U.-G. Meißner and D. Drechsel, *Nucl. Phys. A* 596 (1996) 367
- [231] J. Speth and R. Tegen, *Z. Phys. C* 75 (1997) 717
- [232] C.B. Dover and A. Gal, *Prog. Part. Nucl. Phys.* 12 (1985) 171
- [233] S. Capstick and W. Roberts, *Phys. Rev. D* 49 (1994) 4570
- [234] N. Kaiser, *Phys. Rev. C* 60 (1999) 057001

- [235] A.I. Titov, B. Kämpfer and B.L. Reznik, *Eur. Phys. J. A* 7 (2000) 543
- [236] F. Hibou et al., *Phys. Rev. Lett.* 83 (1999) 492
- [237] N. Kaiser, *Eur. Phys. J. A* 5 (1999) 105
- [238] V. Flaminio et al., *Report CERN-HERA-84-01* (1984)
- [239] V. Herrmann, J. Speth and K. Nakayama, *Phys. Rev. C* 43 (1991) 394
- [240] A.I. Titov, B. Kämpfer and B.L. Reznik, *Phys. Rev. C* 65 (2002) 065202
- [241] D.O. Riska and G.E. Brown, *Nucl. Phys. A* 679 (2001) 577
- [242] J. Ellis, M. Karliner, D.E. Khazeev and M.G. Sapozhnikov, *Nucl. Phys. A* 673 (2000) 256
- [243] R. Jaffe, *Phys. Rev. D* 15 (1977) 267
- [244] F. Close, *Rep. Prog. Phys.* 51 (1988) 833
- [245] F. Close et al., *Phys. Lett. B* 319 (1993) 291
- [246] D. Morgan and M.R. Pennington, *Phys. Rev. D* 48 (1993) 1185
- [247] F. Kleefeld et al., *e-Print Archive: hep-ph/0109158*
- [248] J. Weinstein and N. Isgur, *Phys. Rev. D* 41 (1990) 2236
- [249] Z.S. Wang, S. Krewald and J. Speth, *Nucl. Phys. A* 684 (2000) 429c
- [250] K. Hikasa et al. (Particle Data Group), *Phys. Rev. D* 45 (1992) S1
- [251] D. Lohse et al., *Nucl. Phys. A* 516 (1990) 513
- [252] O. Krehl, R. Rapp and J. Speth, *Phys. Lett. B* 390 (1997) 23
- [253] J. Haidenbauer, Proc. of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 225
- [254] J.A. Oller, E. Oset and J.R. Peláez, *Phys. Rev. D* 59 (1999) 074001
- [255] B.S. Zou and D.V. Bugg, *Phys. Rev. D* 48 (1993) R3948
- [256] N.A. Törnqvist, *Phys. Rev. Lett.* 49 (1982) 624
- [257] R. Kamiński, L. Leśniak and B. Loiseau, *Phys. Lett. B* 413 (1997) 130
- [258] R. Kamiński, L. Leśniak and B. Loiseau, *Eur. Phys. J. C* 9 (1999) 141
- [259] R. Kamiński, L. Leśniak and B. Loiseau, *Acta Phys. Pol. B* 31 (2000) 2467
- [260] J.A. Oller, E. Oset and A. Ramos, *Prog. Part. Nucl. Phys.* 45 (2000) 157
- [261] D. Alde et al., *Z. Phys. C* 66 (1995) 375 , *Phys. Lett. B* 397 (1997) 350 , *Eur. Phys. J. A* 3 (1998) 361
- [262] D. Barberis et al., *Phys. Lett. B* 462 (1999) 462
- [263] D. Barberis et al., *Phys. Lett. B* 453 (1999) 325
- [264] A. Breakstone et al., *Z. Phys. C* 31 (1986) 185
- [265] E.M. Aitala et al., *Phys. Rev. Lett.* 86 (2001) 765 , *Phys. Rev. Lett.* 86 (2001) 770
- [266] R.R. Akhmetshin et al., *Phys. Lett. B* 462 (1999) 371
- [267] R.R. Akhmetshin et al., *Phys. Lett. B* 462 (1999) 380
- [268] K. Ackerstaff et al., *Eur. Phys. J. C* 4 (1998) 19
- [269] P. Abreu et al., *Phys. Lett. B* 449 (1999) 364
- [270] K. Ackerstaff et al., *Eur. Phys. J. C* 5 (1998) 411
- [271] J.B. Gay et al., *Phys. Lett. B* 63 (1976) 220
- [272] S. Teige et al., *Phys. Rev. D* 59 (1998) 012001

- [273] C. Quentmeier, *Ph.D. thesis*, University of Münster, Germany (2001)
- [274] E.L. Bratkovskaya et al., *Eur. Phys. J. A* 4 (1999) 165
- [275] V.Y. Grishina et al., *Eur. Phys. J. A* 9 (2000) 277
- [276] V. Chernyshev et al., COSY Proposal No. 55 (1997), M. Büscher et al., COSY Proposal No. 55.2 (2001); <http://www.fz-juelich.de/ikp/anke>
- [277] E. Oset, J.A. Oller, and U.-G. Meißner, *Eur. Phys. J. A* 12 (2001) 435
- [278] P. Moskal and W. Oelert, Proceedings of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 212
- [279] W. Oelert, Proposal for the COSY-11 facility, COSY (1988)
- [280] M. Büscher et al., COSY Proposal No. 97 (2001) <http://www.fz-juelich.de/ikp/anke>
- [281] S. Steininger and U.-G. Meißner, *Phys. Lett. B* 391 (1997) 446
- [282] U.-G. Meißner, *e-Print Archive*: hep-ph/0108133, hep-ph/0201078 and references therein
- [283] M. Alberg, *Prog. Part. Nucl. Phys.* 36 (1996) 217
- [284] J. Aichelin and C.M. Ko, *Phys. Rev. Lett.* 55 (1985) 2661 , W. Cassing, V. Metag, U. Mosel and K. Niita, *Phys. Rep.* 188 (1990) 363 , U. Mosel, *Ann. Rev. Nucl. Part. Sci* 41 (1991) 29 , G.E. Brown, C.M. Ko, Z.G. Wu and L.H. Xia, *Phys. Rev. C* 43 (1991) 1881 , G.Q. Li, A. Faessler and S.W. Huang, *Prog. Part. Nucl. Phys.* 31 (1993) 159 , B.A. Li, *Phys. Rev. C* 50 (1994) 2144 , G. Hartnack, J. Jaenicke, L. Sehn, H. Stöcker and J. Aichelin, *Nucl. Phys. A* 580 (1994) 643 , G.Q. Li and C.M. Ko, *Phys. Lett. B* 349 (1995) 405
- [285] X.S. Fang, C.M. Ko, G.Q. Li and Y.M. Zheng, *Phys. Rev. C* 49 (1994) 608
- [286] M. Lutz, *Phys. Lett. B* 426 (1998) 12
- [287] A. Ramos, and E. Oset, *Nucl. Phys. A* 671 (2000) 481
- [288] D.B. Kaplan and A.E. Nelson, *Phys. Lett. B* 175 (1986) 57
- [289] G.E. Brown, C.H. Lee, M. Rho and V. Thorsson, *Nucl. Phys. A* 567 (1994) 937
- [290] W. Cassing and E.L. Bratkovskaya, *Phys. Rep.* 308 (1999) 65
- [291] F. Laue et al., *Phys. Rev. Lett.* 82 (1999) 1640
- [292] P. Senger, *Nucl. Phys. A* 685 (2001) 312c and references therein
- [293] H. Oeschler, *J. Phys. G* 27 (2001) 257 and references therein
- [294] B. Schürmann and W. Zwermann, *Phys. Lett. B* 183 (1987) 31
- [295] W. Zwermann, *Mod. Phys. Lett. A* 3 (1988) 251
- [296] J. Randrup and C.M. Ko, *Nucl. Phys. A* 343 (1980) 519 , *Nucl. Phys. A* 411 (1983) 537
- [297] G.Q. Li and C.M. Ko, *Nucl. Phys. A* 594 (1995) 439
- [298] A. Sibirtsev, *Phys. Lett. B* 359 (1995) 29
- [299] K. Tsushima, A. Sibirtsev and A.W. Thomas, *Phys. Lett. B* 390 (1997) 29
- [300] G. Fäldt and C. Wilkin, *Z. Phys. A* 357 (1997) 241
- [301] M. Hoffmann et al., *Nucl. Phys. A* 593 (1995) 341
- [302] A. Deloff, *Nucl. Phys. A* 505 (1989) 583
- [303] M.M. Nagels et al., *Phys. Rev. D* 20 (1979) 1633
- [304] B. Holzenkamp et al., *Nucl. Phys. A* 500 (1989) 485
- [305] A. Reuber et al., *Nucl. Phys. A* 570 (1994) 543
- [306] G. Alexander et al., *Phys. Rev.* 173 (1968) 1452

- [307] B. Sechi-Zorn et al., *Phys. Rev.* 175 (1968) 1735
- [308] E. Ferrari, *Phys. Rev.* 120 (1960) 988 , *Nuovo Cimento* 15 (1960) 652
- [309] D.H. Saxon et al., *Nucl. Phys.* B 162 (1980) 522 , K.W. Ball et al., *Nucl. Phys.* B 222 (1983) 389, D.J. Candlin et al., *Nucl. Phys.* B 226 (1983) 1 , *Nucl. Phys.* B 238 (1984) 477
- [310] J.-M. Laget, *Phys. Lett.* B 259 (1991) 24
- [311] K. Tsushima, S.W. Huang and A. Faessler, *Phys. Lett.* B 337 (1994) 245 , *J. Phys.* G 21 (1995) 33
- [312] G. Giacometti et al., *Nucl. Phys.* B 20 (1970) 301 , *Nucl. Phys.* B 71 (1974) 138
- [313] B.R. Martin, *Nucl. Phys.* B 94 (1975) 413
- [314] P.M.M. Maessen, T.A. Rijken and J.J. de Swart, *Phys. Rev.* C 40 (1989) 2226
- [315] K. Tsushima, A. Sibirtsev, A.W. Thomas and G.Q. Li, *Phys. Rev.* C 59 (1999) 369
- [316] R. Shyam, *Phys. Rev.* C 60 (1999) 055213
- [317] F. Balestra et al., *Phys. Rev. Lett.* 83 (1999) 1534
- [318] S.E. Vigdor, Proceedings of the *International Workshop On Flavor And Spin In Hadronic And Electromagnetic Interactions*, September 21–23, 1992, Turin, Italy; Eds. F. Balestra, R. Bertini, R. Garfagnini (Ed. Compositori, Bologna, 1993), p.317
- [319] A. Bravar et al., *Phys. Rev. Lett.* 78 (1997) 4003
- [320] Z. Liang and C. Boros, *Phys. Rev. Lett.* 79 (1997) 3608
- [321] M. Maggiora et al., *Nucl. Phys.* A 691 (2001) 329
- [322] K. Heller, Proceedings of the *12th International Symp. on High-energy Spin Physics (SPIN 96)*, September 10–14, 1996, Amsterdam, Netherlands; Eds. C.W. de Jager, T.J. Ketel, P.J. Mulders, J.E.J. Oberski, M. Oskam–Tamboezer (World Scientific, Singapore, 1997), S. Paul, *Int. Workshop on Strange Structure Of The Nucleon*, March 11–15, 1997, Geneva, Switzerland, J. Felix, *Mod. Phys. Lett.* A 12 (1997) 363 , *Mod. Phys. Lett.* A 14 (1999) 827
- [323] A. Sibirtsev, K. Tsushima and A.W. Thomas, *Phys. Lett.* B 421 (1998) 59
- [324] A. Sibirtsev, K. Tsushima, W. Cassing and A.W. Thomas, *Nucl. Phys.* A 646 (1999) 427
- [325] N.G. Kelkar, B.K. Jain, *Int. J. Mod. Phys.* E 9 (2000) 431
- [326] R. Siebert et al., *Nucl. Phys.* A 567 (1994) 819
- [327] T.H. Tan, *Phys. Rev. Lett.* 23 (1969) 395
- [328] J.T. Balewski et al., *Eur. Phys. J.* A 2 (1998) 99
- [329] A. Gasparyan *diploma thesis*, Moscow State Engineering Physics Institute, 1999
- [330] A. Gasparian et al., *Phys. Lett.* B 480 (2000) 273
- [331] R.H. Dalitz, J. McGinley, C. Belyea and S. Anthony, Proceedings of the *Int. Conf. on Hypernuclear and Kaon Physics*, 1982, Heidelberg, Germany; Ed. B. Povh, Report MPI H-82-V20 (Max-Planck-Institut Heidelberg, 1982)
- [332] A. Martin, *Nucl. Phys.* B 179 (1981) 33
- [333] J.J. de Swart, *Rev. Mod. Phys.* 35 (1963) 916
- [334] D. Cline, R. Laumann and J. Mapp, *Phys. Rev. Lett.* 20 (1968) 1452
- [335] T.H. Tan, *Phys. Rev.* D 7 (1973) 600
- [336] A. Kudryavtsev, *JETP Lett.* 14 (1971) 90
- [337] R.H. Dalitz and A. Deloff, *Czech. J. Phys.* B 32 (1982) 1021 , *Aust. J. Phys.* 36 (1983) 617
- [338] E. Oset et al., *Phys. Rep.* 188 (1990) 79

- [339] C.J. Batty et al., *Phys. Lett. B* 74 (1978) 27
- [340] A.M. Gasparian et al., *Nucl. Phys. A* 684 (2001) 397
- [341] A. Sibirtsev, K. Tsushima, W. Cassing and A.W. Thomas, *e-Print Archive* nucl-th/0004022
- [342] R. Shyam, G. Penner and U. Mosel, *Phys. Rev. C* 63 (2001) 022202
- [343] A. Engel, R. Shyam, U. Mosel and A.K. Dutt-Majumder, *Nucl. Phys. A* 603 (1996) 387
- [344] J. Dubach, W.M. Kloet and R.R. Silbar, *Phys. Rev. C* 33 (1986) 373
- [345] L. Tiator, C. Bennhold and S.S. Kamalov, *Nucl. Phys. A* 580 (1994) 455
- [346] J.-M. Laget, *Nucl. Phys. A* 691 (2001) 11c
- [347] A. Gasparian, Proc. of the *Symp. on Threshold Meson Production in pp and pd Interactions*, June 20–24, 2001, Cracow, Poland; Eds. P. Moskal and M. Wolke, *Schriften des FZ-Jülich – Matter and Materials* 11 (2002) 205
- [348] J. Haidenbauer, W. Melnitchouk and J. Speth, Proceedings of the *8th Meeting On Mesons And Light Nuclei*, July 2–6, 2001, Prague, Czech Republic; *AIP Conf. Proc.* 603 (2001) 421
- [349] J. Haidenbauer, private communication
- [350] A.T.M. Aerts and C.B. Dover, *Nucl. Phys. B* 253 (1985) 116
- [351] H. Piekarz, *Nucl. Phys. A* 479 (1988) 263c
- [352] O. Braun et al., *Nucl. Phys. B* 124 (1977) 45
- [353] F. Hinterberger et al., COSY Proposal No. 92 (2000)
- [354] J.A. Gómez Tejedor and E. Oset, *Nucl. Phys. A* 571 (1994) 667
- [355] E. Oset and M.J. Vicente-Vacas, *Nucl. Phys. A* 446 (1985) 584
- [356] L. Alvarez-Ruso et al., *Nucl. Phys. A* 633 (1998) 519
- [357] J. Johanson, *Ph.D. thesis*, Uppsala University (2000)
- [358] W. Brodowski et al., *Phys. Rev. Lett.* 88 (2002) 192301
- [359] D.R. F. Cochran et al., *Phys. Rev. D* 6 (1972) 3085
- [360] F.H. Cverna et al., *Phys. Rev. C* 23 (1981) 1698
- [361] D.V. Bugg et al., *Phys. Rev.* 133 (1964) B1017
- [362] L.G. Dakhno et al., *Sov. J. Nucl. Phys.* 37 (1983) 540
- [363] F. Shimizu et al., *Nucl. Phys. A* 386 (1982) 571
- [364] D.C. Brunt et al., *Phys. Rev.* 187 (1969) 1856
- [365] J. Stepaniak, *VI. Int. Oberjoch Meeting on Meson-Nuclear Physics*, Oberjoch, Germany (1996)
- [366] L. Alvarez-Ruso, *Ph.D. thesis*, University of Valencia (1999)
- [367] A. Abashian, N.E. Booth and K.M. Crowe, *Phys. Rev. Lett.* 5 (1960) 258
- [368] A. Abashian et al., *Phys. Rev.* 132 (1963) 2206
- [369] J. Banaigs et al., *Nucl. Phys. B* 67 (1973) 1
- [370] G. Grayer et al., *Nucl. Phys. B* 75 (1974) 189
- [371] T. Risser et al., *Phys. Lett. B* 43 (1973) 68
- [372] F. Bellemann et al., *Phys. Rev. C* 60 (1999) 061002
- [373] L. Alvarez-Ruso, *Phys. Lett. B* 452 (1999) 207 ;
E. Oset et al., *Few Body Syst.Suppl.* 11 (1999) 275
L.Alvarez-Ruso, E.Oset and E.Hernandez, Proceedings of the *7th International Conference On Mesons And Light Nuclei 98*, August 31 – September 4, 1998, Pruhonice, Prague, Czech Republic; Eds. J. Adam, P. Bydzovsky, J. Dobes, R. Mach, J. Mares, M. Sotona (World Scientific, Singapore, 1999), p.81, *e-Print Archive*:nucl-th/9810002;

- [374] V. Betker et al., *Phys. Rev. Lett.* 77 (1996) 3510
- [375] E. Plouin et al., *Nucl. Phys. A* 302 (1978) 413
- [376] C.L. Hollas et al., *Phys. Rev. C* 25 (1982) 2614
- [377] F. Bonutti et al., *Nucl. Phys. A* 638 (1998) 729
- [378] M.Kermani et al., *Phys. Rev. C* 58 (1998) 3419
- [379] F. Bellemann et al., *Annual Report 2001*, IKP, Berichte des FZ-Jülich, 3978, p44
- [380] J. Ritman et al. *Nucl. Phys. A* 691 (2001) 395
- [381] T. Waas et al., *Phys. Lett. B* 379 (1996) 34 ; T. Waas, N. Kaiser, and W. Weise, *Phys. Lett. B* 365 (1996) 12 ; T. Waas, and W. Weise, *Nucl. Phys. A* 625 (1997) 287
- [382] V. Koch, *Phys. Lett. B* 337 (1994) 7
- [383] A. Baca, C. Garcia-Recio, J. Nieves, *Nucl. Phys. A* 673 (2000) 335
- [384] A. Baldini et al., *Total Cross-Sections for Reactions of High Energy Particles*, Ed. H. Schopper, Landolt-Börnstein, New Series I/12 (Springer, Berlin, 1988)
- [385] J. Berger et al., *Phys. Rev. Lett.* 61 (1988) 919
- [386] B. Mayer et al., *Phys. Rev. C* 53 (1996) 2068
- [387] Q. Haider and L.C. Liu, *Phys. Lett. B* 172 (1986) 257
- [388] L.C. Liu and Q. Haider, *Phys. Rev. C* 34 (1986) 1845
- [389] S. Wycech et al., *Phys. Rev. C* 52 (1995) 544
- [390] V.V. Abaev and B.M.K. Nefkens, *Phys. Rev. C* 53 (1996) 385
- [391] H.C. Chiang, E. Oset and L.C. Liu, *Phys. Rev. C* 44 (1991) 738
- [392] M. Betigeri et al., *Phys. Lett. B* 472 (2000) 267
- [393] R. Bilger et al., *Phys. Rev. C* 65 (2002) 044608
- [394] K. Kilian and H. Nann, *AIP Conf. Proc.* 221 (1990) 185
- [395] G. Fäldt and C. Wilkin, *Nucl. Phys. A* 587 (1995) 769
- [396] *WebElements*²⁰⁰⁰, the periodic table on the WWW, <http://www.webelements.com>,
I. Mills et al., *Quantities, Units and Symbols in Physics Chemistry*, (Blackwell Scientific Publications, Oxford, 1988)
- [397] B. Krusche et al., *Phys. Rev. Lett.* 74 (1995) 3736 , *Phys. Rev. Lett.* 75 (1995) 3023
- [398] G.L. Li, W.K. Cheung and T.T.S. Kuo, *Phys. Lett. B* 195 (1987) 515
- [399] F. Frascaria et al., *Phys. Rev. C* 50 (1994) R537
- [400] N. Willis et al., *Phys. Lett. B* 406 (1997) 14
- [401] H.O. Meyer, *Phys. Scripta* 48 (1993) 54
- [402] J.A. Niskanen, *Acta Phys. Pol. B* 31 (2000) 2683
- [403] M. Betigeri et al., *Nucl. Phys. A* 690 (2001) 473
- [404] P. Hawranek, Proceedings of the 5th International Conference on Nuclear Physics at Storage Rings, STOR1'02, June 16–20, 2002, Uppsala, Sweden
- [405] A. Magiera and H. Machner, *Nucl. Phys. A* 674 (2000) 515
- [406] A. Magiera et al., COSY Proposal No. 59, <http://ikpgem02/gem/gem.html>
- [407] C. Wilkin, *Phys. Lett. B* 331 (1994) 276
- [408] J. Bacelar et al., COSY Proposal No. 83.2 (2000), <http://www.fz-juelich.de/ikp/anke>
- [409] W.R. Lozowski and J.D. Hudson, *Nucl. Instr. and Meth. A* 303 (1991) 34
- [410] V. Jaeckle et al., *Nucl. Instr. and Meth. A* 349 (1994) 15

- [411] A. Hassan et al., *Nucl. Instr. and Meth. A* 425 (1999) 403
- [412] A. Khoukaz et al., *Eur. Phys. J. D* 5 (1999) 275
- [413] B. Trostell, *Nucl. Instr. and Meth. A* 362 (1995) 41
- [414] C. Ekström et al., *Nucl. Instr. and Meth. A* 371 (1996) 572
- [415] C. Ekström et al., *Phys. Scripta* T00 (2002) 1
- [416] F. Sperisen et al., *IUCF Sci. and Techn. Rep. 1989/90* 117
- [417] H. Dombrowski et al., *Nucl. Instr. and Meth. A* 386 (1997) 228 and references therein
- [418] H. Calén et al., *Nucl. Instr. and Meth. A* 379 (1996) 57
- [419] M. Dahmen et al., *Nucl. Instr. and Meth. A* 348 (1994) 97
- [420] G. Anton et al., *Nucl. Instr. and Meth. A* 310 (1991) 631
- [421] W.W. Daehnick et al., *Phys. Rev. C* 65 (2002) 024003 and references therein
- [422] W. Deinet et al., *Nucl. Phys. B* 11 (1969) 495
- [423] C. Wilkin, private communication
- [424] T.-S.H. Lee, C. Hanhart, private communication
- [425] A.B. Santra and B.K. Jain, *Nucl. Phys. A* 634 (1998) 309
- [426] E. Gedalin et al., *Nucl. Phys. A* 634 (1998) 368
- [427] Proceedings of the *Summer School and Workshop on COSY Physics*,
 August 30 – September 4, 2002, Kerkrade, The Netherlands;
 organized by K. Brinkmann, M. Büscher, C. Hanhart, A. Khoukaz, A. Lehrach, F. Rathmann and
 M. Wolke, <http://www.fz-juelich.de/css2002>, to be published.